

species has the facies of *Lygocoris (Apolygus)* Reuter.

***Lygus v-nigrum* Poppius, 1915**

One female, typus, Formosa, Sauter, Chip Chip, 1909.II. Color greenish straw, a V-like black spot on apex of clavus and two V-like dark spots at apex of endocorium, apical portion of antenna II and segments III-IV black. There is a male cotype.

***Macrolonius schenklingi* Poppius, 1915**

Two female specimens (typus and cotypus), Formosa, Sauter, Fuhosho, 1909, VI.

***Macropeplus brunnescens* Poppius, 1912**

One female, typus, New Guinea, Biró, 1897, Stephansort, Austrolabe Bai. Brown testaceous, pronotum and scutellum rugose. Size very large, elongate, abdomen, head and propleura pumpkin color, rostrum reaching the middle coxae.

***Megacoelum minutum* Poppius, 1915**

One male, typus, Formosa, Sauter, Takan, 1907. Size small, segment I of antenna with reddish dots, neck red, abdomen and body below marmorate with reddish and brown.

***Orthotylus orientalis* Poppius, 1915**

One specimen, male, typus, Formosa, Sauter, Amping, 1908, VI. Small, greenish, head, pronotum and legs ochraceous, membrane fuscous. It belongs to subgenus *Melanotrichus* Reuter, 1875.

***Pachypeltis annulipes* Poppius, 1912**

One female, typus, New Guinea, Biró, 1899, Simbang, Huon-Golf. Specimen mutilated. Head and antenna black (segment I lighter), abdomen with segments II-IV milky white and segments IV-VIII with white spots, femur III with a pale ring, membrane with three small red spots, cuneus very long, apex of areola reaching much beyond apex of cuneus.

***Pachypeltis politum* (Walker, 1873)  
Poppius, 1912**

One specimen, Borneo. Brown with two black spots at apex of hemelytra (beyond apex of clavus). Size large.

***Pachypeltopsis australicus* Poppius, 1912**

One female, Australia, N.S. Wales. Shining, brown, basal third of corium pale, apical half of hemelytra black, calli fused with a depression behind, glabrous (?), segments II-IV of abdomen white, segments V-VIII with white spots. Size large.

***Periscopos mundulus* (Breddin, 1896)  
Carvalho & Southwood, 1955**

One female; typus, Java. Head, pronotum, scutellum, antenna (except segment I) black, middle of hemelytra infuscate, membrane short. Two males cotypes, both with two pale spots on vertex.

***Peritropis annulicornis* Poppius, 1909**

Four specimens, New Guinea, Simbang, Huon-Golf. Similar to those in the Museum of Natural History "Giacomo Doria", Genova.

***Peritropis lugubris* Poppius, 1909**

The typus is lost. Only pin and label remains. Singapore, Biró, 1898.

***Pilophorus formosanus* Poppius, 1914**

One female, typus, Formosa, Sauter, Kosempo, 1908. Very small, four lines and humeral angles silvery; hemelytra beyond apex of clavus, apex of second antennal segment and scutellum black, head, pronotum dark brown, the basal half of hemelytra brown.

***Platypeltocoris planus* Poppius, 1912**

One female, typus, New Guinea, Biró, 1899, Sattelberg, Huon-Golf. Black, base of first antennal segment and legs pale. Size medium.