

Platypeltocoris similis Poppius, 1912

One male, *typus*, New Guinea, Biró, 1900, Friedrich Wilhelm Haffen. Pygophore with left paramere sclerotized and pipe-like in shape.

Pleurochilophorus subhyalinus Poppius, 1912

One female, *typus*, Camerum, Adawana. Transparent, two small black spots at humeral angles, claval commissure reddish.

Proboscidoecoris neoguineanus Poppius, 1915

One male, *typus*, New Guinea, Biró, 1897, Erima, Astrolabe Bai. Size small, black, hind margin of disc, apex of scutellum and corium contiguous to cuneal fracture pale, antenna pale, apex of segment II and segments III-IV black, apex of segment I of rostrum, xyphus, lower margin of propleura and lower half of ostiolar peritreme white.

Proboscidoecoris taivanus Poppius, 1915

One male, *typus*, Formosa, Sauter, Tainhorinsho, 1909, VIII. Black, size small; spots on vertex near eyes, base of endocorium, apices of clavus and scutellum, cuneal fracture, base and apex of cuneus pale; hind femur with three pale fasciae.

Prodromopsis basalis Poppius, 1915

Two females, *cotypes*, Formosa, Sauter, Tatherin, I.1910. Pale, eyes, segments II-IV of antenna black, cuneus more or less curved externally.

Prodromopsis cuneatus (Distant, 1909)
Poppius, 1911

One male, Ceylon, Madarasz, Kala-Wewa, 1896, Feb.10. Pronotal disc and hemelytra more or less lutescent, second antennal segment reddish at apex and base.

Prodromopsis oculatus Poppius, 1911

One male, *typus*, New Guinea, Biró, Erima, Austrolabe Bai. Pale with head and pronotum lutescent, claval commissure reddish, left paramere

large and curved upwards, black; antenna black (except narrowed base of segment I); eyes strongly pedunculate upwards.

Prodromopsis scutellaris Poppius, 1914

One female, *typus*, Formosa, Sauter, Tathorin, 1910, V. Pale, segments II-IV of antenna, hind margin of disc, scutellum and eyes black.

Pycnofurius pallidiscutum Poppius, 1915

One female, *typus*, Formosa, Sauter, Pilam, 1908. Pale, head, pronotum, scutellum, clavus, two roundish spots on corial commissure (one at each side), areola and apex of cuneus black; legs and coxae pale; apical half of second antennal segment and segments III-IV black.

Rhinocylapidius velocipedoides Poppius, 1915

Three specimens, *typus* female, Formosa, Sauter, Huhosho, IV.1909. Head horizontal, very long, pronotum with a T-like pale spot behind calli, membrane short, embolium very wide and punctate, rostrum reaching beyond apex of pygophore.

Rhodolygus obscuritarsis Poppius, 1915

One male, *typus*, Australia, Biró, 1900, Sydney, N.S. Wales. Ochraceous reddish: a longitudinal fascia and oblique striae on vertex, neck, hind portion of disc, scutellum (except longitudinal median fascia), mesoscutum (except two black spots), clavus (except apex), endocorium on apical portion, veins of membrane, apical portion of third femur and base of third tibia, upper surface and longitudinal spots on abdomen and antenna fuscous.

Stenotus insularis Poppius, 1915

One female, *typus*, Formosa, Sauter, Amping, 1908. Very small, straw color, head pointed, membrane fuscous.

Stenotus elegans Poppius, 1912

Two females, *typus*, Deutsch O. Afrika. Head and pronotum lutescent, hemelytra citrine