

with endoclavus, inner margin of endocorium, oblique fascia at middle of corium, paracuneus, membrane and antenna black, legs brown.

***Stenotus longiceps* Poppius, 1915**

One specimen, Formosa, Sauter, Amping, 1908. Small, first antennal segment, apical spot on endocorium and dots on inner margin of cuneus reddish, femora with red spots.

***Stenotus pygmaeus* Poppius, 1915**

One male, typus, Formosa, Sauter, Amping, 1906. Very small, citrine, two longitudinal fasciae on pronotum, internal margins of eyes, scutellum, clavus and two apical spots on endocorium, membrane black.

***Taedia notaticollis* (Reuter, 1907)  
Carvalho, 1959**

Two specimens, male typus, Paraguay, Vezenyi. Size medium, two elongate spots behind calli black, scutellum pale at sides, brown at middle with a longitudinal pale fascia, propleura with three longitudinal brown fasciae.

***Taedia trivittata* (Reuter, 1913)  
Carvalho, 1959**

Two specimens, females, typus and cotypus, Peru, Marcapata. Size large, elongate, black, calli, longitudinal median fascia on disc, middle of scutellum, claval commissure, longitudinal fascia on exocorium, basal half of embolium and cuneus lutescent.

***Thaumastomiris discoidalis* Poppius, 1912**

One male, typus, New Guinea, Biró, 1896, Seleó, Berlinhaf. Red with a dark spot on clavus and endocorium.

***Thaumastomiris sanguinalis* Kirkaldy, 1902**

Two males, typus and cotypus, Ceylon, Peradneya. Red, eyes, antenna and claws black, membrane fuscous.

***Tinginotum camelus* Poppius, 1915**

One female, typus, New Guinea, Biró, 1899, Sattelberg, Huon-Golf plus two cotypes. Pronotum with a prominent tubercle at middle of disc (*camelus*), hemelytra with pruinose silvery spots, a transversal fascia on hemelytra and clavus black, apex of third femur with three rings and third tibia also with three rings black.

***Tinginotum cretaceum* Poppius, 1915**

One male, typus, New Guinea, Biró, 1898, Sattelberg, Huon-Golf. Pronotum without a prominent tubercle but with tufts of hairs, embolium with 6-7 fasciae or spots dark, second antennal segment with four pale spots. It is a synonym of *Tinginotum papuanum* Poppius, 1915.

***Tinginotum formosanum* Poppius, 1915**

One female, typus, one male, cotypus, Formosa, Sauter, Mt. Hoozan 1909, XII. Embolium with 8-9 spots or fasciae, second antennal segment with two pale fasciae.

***Tinginotum papuanum* Poppius, 1915**

One female, typus, New Guinea, Biró, 1896, Erima, Astrolabe Bai. Large, pronotum with tufts of hairs, scutellum with two black spots, first antennal segment dark apically and an oblique dark line at base, embolium wide, well marked with brown cross bars. This species is described from a female of *Tinginotum cretaceum* Poppius and has page priority over it.

***Trilaccus nigroruber* Horvath, 1902**

Three specimens, male, typus, Australia, N.S. Wales, Biró. Black, metallic, head and pronotum (except hind margin), corium beyond apex of clavus red, calli fused, transversal, with a fossa behind them.

***Tyraquellus brunneus* Poppius, 1915**

One male, typus, Formosa, Sauter. This species was transferred to genus *Hallodapus* Fieber by Carvalho (1958).