

col., 90-115. The species do not pertain to genus *Argenis* Distant and may constitute a new genus.

***Argenis incisuratus* (Walker, 1873)**

The type is a female from Thwaites, Ceylon.

***Bothriomiris dissimulans* (Walker, 1873)**

The scutellum is reddish with a pale longitudinal fascia.

***Bothriomiris testaceus* Distant, 1904**

Color yellowish testaceous, membrane fuscous.

***Carmelus fasciatus* Distant, 1893**

This species shows a dark transversal fascia on corium level with apex of scutellum.

***Carmelus formosus* (Distant, 1884)**

With a dark spot in the middle of clavus, embolium, apex of exocorium and margins of cuneus reddish.

***Carmelus lunatus* (Distant, 1884)**

A spot on disc posteriorly, base of scutellum and a spot at middle of clavus, apex of corial commissure and membrane black, hind margin of pronotum, clavus, transversal fascia on corium and embolium level with apex of clavus reddish.

***Carmelus parvus* Distant, 1884**

Reddish ochraceous, spots behind calli, humeral angles, scutellum and embolium at base white.

***Clapmarius turgidus* Distant, 1904**

The first antennal segment as in *Lampethusa* Distant.

***Cylapofulvius listeri* Izzard, 1936**

This species from Christmas Islands has the pronotum and scutellum punctate, hemelytra

shagreen, smooth, cuneal suture small, oblique, embolium widening to base, color brown mottled with pale. Its position in the genus is questioned.

***Cyrtocapsus marginatus* (Distant, 1893)**

Black with pale embolium, legs black with base of femora and apical 3/4 of tibiae pale, antenna ochraceous.

***Derophthalma emissitia* (Distant, 1893)**

Lectotype female, Orizaba, with segment I of antenna black only at base. Paralectotype female, from Rio Honda, Honduras, with segment I of antenna black, pale apically.

***Derophthalma laterata* (Distant, 1883)**

Lectotype female and two paralectotypes from Chiriqui, Panama. Segment I of antenna black.

***Dichroocoris nigripes* (Distant, 1884)**

Black, head, pronotum, mesoscutum and first rostral segment red; size small.

***Eccritotarsus argentatus* (Distant, 1884)**

Ochraceous with four rounded black spots (two at base of clavus and two at middle of corium), tibiae and apex of cuneus black.

***Eccritotarsus linearis* (Distant, 1884)**

Ochraceous with longitudinal black fasciae on body (two at middle of corium and two covering embolium), margins of cuneus and nervures of membrane black.

***Eccritotarsus pictus* (Distant, 1884)**

Ochraceous, scutellum, base and apex of clavus, a transversal fascia on corium level with apex of clavus black (these fasciae widen inwards), apex of second antennal segment and third tarsi black.