

Orthotylus (Melanotrichus) flavosparsus (C. Sahlberg 1842)

Aralik (Horváth 1894 and Kiritschenko 1918).

Species of Holarctic distribution, in S WAsia recorded from Caucasia and Transcaucasia, in Central Asia from Turkestan.

Orthotylus (Melanotrichus) beieri E. Wagner 1942

2 ♂♂ and 1 ♀ — Toros: Karapinar, 1. VIII. 1947. (Exp. of the National Museum, Praha.)

On the undergrowth of light, dry forest among *Juniperus excelsa* and *Abies cilicica*.

These specimens constitute a new record for Turkey.

This species up till now recorded only from the type locality in South Carinthia, and from S. France.

Orthotylus (Orthotylus) cupressi Reuter 1883

1 ♀ — Hasanoğlan, 13. VII. 1947. (Exp. of the National Museum, Praha.)

Flying at dusk high above the ground in the region of wild fruit-trees at an altitude of 1.000 m.

This specimen constitutes a new record for Turkey.

Species so far known from the western Mediterranean.

Orthotylus (Orthotylus) marginalis Reuter 1884

2 ♂♂ — Edirne, 8.—13. VI. 1947. (Exp. of the National Museum, Praha.)

On the growth of *Ulmus*, along the river Meriç.

These specimens constitute a new record for Turkey.

Species of Eurosiberian distribution, in SW Asia recorded from Caucasia and Transcaucasia.

Orthotylus (Orthotylus) virescens (Douglas and Scott 1865)

Asia Minor (Oshinin 1909, 1912).

Species of Holomediterranean distribution, extending northwards as far as England. In the Middle East recorded only from Asia Minor and Cyprus.

Brachynotocoris vitcinus Seidenstücker 1954

2 ♂♂ and 3 ♀♀ — Toros: Kozan, 8.—9. VIII. 1947. (Exp. of the National Museum, Praha.)

In the deciduous forest formation.

These specimens constitute a new record for Turkey. This species is described from Syria, NW of Hama (type locality), occurring on *Vitex agnus castus* L. (Seidenstücker).