

**Megalocoleus aurantiacus** (Fieber 1858)

Asia Minor (Reuter 1879 b).

Species of Holomediterranean distribution; Asia Minor is only listed record of the distribution in the Middle East.

**Megalocoleus molliculus** (Fallen 1829)

3 ♂♂ and 2 ♀♀ — Ankara-Baraj, 3.—4. VII. 1947. (Exp. of the National Museum, Praha.)

Occurring on the xerophilous vegetation in the border zone of cultivated area.

Up till now this species has been recorded in Turkey only from Eylar (Horváth 1899, Kiritschenko 1918).

Species of European distribution, in SW Asia recorded from Caucasia and from Central Asia (Turkestan).

**Megalocoleus krueperi** (Reuter 1879)

Asia Minor (Oshinin 1909, 1912).

Probably species of Holomediterranean distribution, recorded also from Algeria, Sicily and Syria.

**Amblytylus concolor** Jakovlev 1877

4 ♀♀ — Beynam, 28. VI. 1947. (Exp. of the National Museum, Praha.)

On Gramineae in wooded hills formation.

Up till now this species has been recorded in Turkey only from Karaköy-Bozüyük (Horváth 1905 a).

Species of Pontomediterranean distribution, in the Middle East recorded only from Central Anatolia, in Central Asia from Tajikistan.

**Amblytylus nasutus** (Kirschbaum 1855)

1 ♂ — Edirne, 8.—13. VI. 1947. (Exp. of the National Museum, Praha.)

Occurring in the border zone of steppe formation.

This specimen constitutes a new record for Turkey.

Species of Mediterranean distribution, penetrating into Central and West Europe. In SW Asia listed only from Transcaucasia.

**Macrotylus quadrilineatus** (Schrank 1785)

Asia Minor (Oshinin 1909, 1912).

Species of Holomediterranean distribution, in the Middle East recorded from Cyprus and Asia Minor.