

**Macrotylus herrichi (Reuter 1873)**

2 ♂♂ and 4 ♀♀ — Ankara-Baraj, 3.—4. VII. 1947. (Exp. of the National Museum, Praha.)

On low vegetation in the border zone of cultivated area.

In Turkey recorded only from Konya-Kayseri (Horváth 1905 a).

Species of Holomediterranean distribution; the Anatolian locality is the only known one in the Middle East.

**Macrotylus seidenstückeri E. Wagner 1954**

Toros: Namrun (E. Wagner 1954 a). Occurring on *Salvia grandiflora* in *Abies* and *Cupressus* forest.

So far known only from the above-mentioned type locality.

**Macrotylus antennalis Horváth 1905**

Yilanli dağ near Kayseri (Horváth 1905 a).

Species up till now recorded only from the above mentioned type locality; probably an endemic of Central Anatolian plateau.

**Macrotylus bipunctatus Reuter 1889**

Asia Minor (Oshanin 1909, 1912).

Species of Holomediterranean distribution; in the Middle East recorded only from Asia Minor.

**Camptotylus linae (Puton 1881)**

Aralik (Horváth 1894, Kiritschenko 1918).

Probably species of Iranian distribution, listed up till now only from Caucasia, Syria and Israel.

**Camptotylus reuteri Jakovlev 1881**

20 ♂♂ and 14 ♀♀ — Toros: Feke, 12. VIII. 1947. (Exp. of the National Museum, Praha.)

On *Tamarix* sp. on the sandy bank of the river Göksu.

These specimens constitute a new record for Turkey.

Species of Pontomediterranean distribution, in the Middle East recorded from Caucasia, Transcaucasia and Cyprus.

**Harpocera thoracica (Fallen 1807)**

Turkey (Thrace) (Oshanin 1909).

Species of Holomediterranean distribution, extending northwards as far as England. In the Middle East recorded also from Caucasia and North Iran.