

Harpocera hellenica Reuter 1876

Bursa (Horváth 1883 a, Reuter 1883); Anatolia (Reuter 1890 a).

Species of Pontomediterranean distribution, recorded also from Greece, Syria and Israel.

Orthonotus rufifrons (Fallen 1807)

Izmir (Reuter 1878) and Gök dağ (Fahringer 1922), occurring on *Origanum vulgare*.

Species of Holomediterranean distribution, extending northwards as far as South Finland and Sweden.

Orthonotus rossicus (Reuter 1878)

Tarsus, Izmir (Reuter 1904 b, 1913).

Species of Pontomediterranean distribution; the two Anatolian localities constitute the only record from the Middle East.

Orthonotus rossicus var. fraudatrix (Reuter 1904)

Tarsus (Reuter 1904 b, 1913)

Variety only known from the above-mentioned type locality.

Phylus (Gnostus) breviceps Reuter 1899

Gaziantep (Reuter 1899).

Species known only from the above mentioned type locality; it seems to be a species of endemic distribution restricted to the northern part of the Syrian plateau.

Ephippicoris lunatus Poppius 1912

Aralik (Kiritshenko 1918).

Probably species of Irano-Turanian distribution, up till now recorded only from Turkestan, Transcaucasia and easternmost part of Turkey.

Psallus (Apocremnus) kolenatii (Flor 1860)

Eğnaki (Kiritshenko 1918).

Species of European distribution, restricted to the zone of pineforests. Eğnaki is the only record in the distribution of this species from SW Asia, and, as N. A. Kiritshenko states it needs reexamination.