

Sthenarus (Phoenicocoris) quercicola Reuter 1904

Menemen (Gediz) (Reuter 1904 b).

Known only from the above-mentioned type locality; probably species of the distribution restricted to Aegean part of Anatolia.

Auchenocrepis reuteri Jakovlev 1876

8 ♂♂ and 7 ♀♀ — Ankara-Baraj, 3.—4. VII 1947. (Exp. of the National Museum, Praha).

On *Tamarix* sp. in the border zone of cultivated areas.

This species has been recorded so far in Turkey from Izmir (Reuter 1878), Aralik (Horváth 1894, Kiritschenko 1918).

Species of Pontomediterranean distribution, in SW Asia recorded from Caucasia, Transcaucasia, Iran and N. Iraq, in Central Asia recorded from Turcomania and Turkestan.

Maurodactylus alutaceus var. discifer Reuter 1901

Amasya (Reuter 1901 b).

Probably species of Holomediterranean distribution, up till now recorded from Spain, variety *discifer* Reuter recorded from NW Anatolia and Turkomania.

Maurodactylus albidus (Kolenati 1845)

3 ♂♂ and 10 ♀♀ — Moğan gölü, 5. VII. 1947. (Exp. of the National Museum, Praha.)

On the low vegetation of steppe formation.

Up till now has been recorded in Turkey only from Aralik (Horváth 1894), Burulan (Kiritschenko 1918).

Species of Caspian distribution, in SW Asia recorded from Caucasia and Transcaucasia, in Central Asia from Turkestan.

Maurodactylus albidus var. discoidalis Reuter 1902

Amasya (Reuter 1902).

Variety recorded only from the above mentioned type-locality.

Tuponia eckerleini E. Wagner 1955

5 ♂♂ and 1 ♀ — Edirne, 8.—13. VI. 1947. (Exp. of the National Museum, Praha.)

On shrubs in the border zone of steppe formation.

This species is described from Mediterranean (type locality: Au Champ du Bés).

Up till now there is recorded from Turkey *Tuponia tamaricis* (Perris 1857) only from Aralik (Horváth 1894 and Kirithenko 1918). This record belongs probably for an another species.