

### **Psallus salicellus** (Mey. D., 1843)

1 ♂ — Slovakia: Zadiel-plateau, 28. VIII. 1959, on *Corylus avellana*, collected Hoberlandt.

Species recorded in Czechoslovakia only from single localities — Bohemia: Jindřichův Hradec, Teplice, Stráž pod Ralskem (Duda, 1885, 1892; Scholtz, 1930), Moravia: Brno, Kuřim, Zbrašov near Hranice, Ondřejov, Široká Niva (Stehlík, 1961), Slovakia: Bardějov, Slovenský kras (Balthasar, 1937). Species of European distribution.

### **Atractotomus tigripes** (Muls., 1852)

39 ♂♂ and 13 ♀♀ — Slovakia: Kovačovské kopce (hills), in steppe conditions, collected Hoberlandt.

Up to the present time only few records are known from Czechoslovakia — Moravia: Mohelno, Březník (Stehlík, 1944), Kobyly (Hoberlandt, 1947, Stehlík, 1948), Javorník (Hoberlandt, 1947), Slovakia: Zadiel, Súľovské skály (Stehlík & Hoberlandt, 1954). Species of European distribution with rather southern extension.

### **Hallodapus suturalis** (H. Sch., 1839)

55 ♂♂ and 32 ♀♀, 10 nymphs — Slovakia: Čenkov, 7. VI. 1960, in sandy conditions, collected Hoberlandt.

Only two records from Czechoslovakia are so far known — Moravia: Pouzdřany (Stehlík & Hoberlandt, 1954), Slovakia: Somotor (Balthasar, 1937). Species of Mediterranean distribution, in Central Europe recorded from Hungary only.

### **Systemonotus kiritshenkoi** Poppius, 1912

*Systemonotus kiritschenkoi* Poppius, 1912, Rev. Russ. d'Ent., 12: 202—203.

*Systemonotus kiritschenkoi*; Kiritschenko, 1912, Rev. Russ. d'Ent., 12: 364.

*Systemonotus kormilevi* Horváth, Kormilev, 1936, Bull. Soc. Sc. Skoplje, 17: 49 and 54. Nomen nudum.

*Systemonotus skopljensis* Kormilev, 1939. Bull. Soc. Sc. Skoplje, 20: 197. **New synonymy.**

*Systemonotus skopljensis*; Wagner, 1947, Entom. Monthly Mag., 84: 4 and 5, fig. 2 B.

*Systemonotus kiritschenkoi*; Kiritschenko, 1951, Nastojaščie polužestokokrylye evrop.

časti SSSR. Opred. po faune SSSR, 42: 171.

*Systemonotus kiritschenkoi*; Stichel, 1956, I. Bestimmungstabellen der Wanzen, II. Europa. 1: 400 and 825.

*Systemonotus skopljensis*; Stichel, 1956, III. Bestimmungstabellen der Wanzen, II. Europa. 1: 401 and 825.

*Systemonotus skopljensis*; Josifov, 1958, Izvestija na zool. inst., 7: 346.

*Systemonotus kiritschenkoi*; Carvalho, 1958, Arq. do Mus. Nac., 45: 178.

*Systemonotus skopljensis* (*scopiensis*); Carvalho, 1958, Arq. do Mus. Nac., 45: 178.

*Systemonotus skopljensis*; Josifov, 1960, Izvestija na zool. inst., 9: 137.

Species originally described by Poppius (1912) from Crimea (Kerch as type locality) and later again recorded from Crimea, Kerch by Kiritschenko (1912). Kormilev (1939) described *Systemonotus skopljensis* from Skopje and Donje Nerezi in S. Yugoslavia, when formerly (Kormilev, 1936) recorded under the unpublished name of *S. kormilevi* Horváth (nomen nudum). Kiritschenko (1951) records this spe-