

*W.* Deer Lake, 17—19. VIII. 49, 2 ind., 24. VII. 51, 3 ind. (L).

*E c o l o g y*: On the bottom on open sandy localities.

*D i s t r i b u t i o n*: Nearctic species with northern distribution. From Newfoundland, Nova Scotia and Quebec to British Columbia. Only in the northern States.

*Nabis* Latr.

*N. flavomarginatus* Schltz

*Nabis flavomarginatus* Schltz, Arb. Schles. Ges. Vat. Kult. 1846, 114. — v. D. 280. — Bl. 600. — Torre Bueno, Cand. Ent. LXII, 6. — Walley, Can. Ent. LXII, 76. — Mo. 16. — Strickl. 199.

*S.E.* Labrador: Forteau, 12—15. VII. 51, 3 ind. (L).

*S.E.* St. Johns, 30. VIII. 49, 1 ind. (L); Come-by-Chance, 27. VIII. 49, 1 ind. (L). — S. Terrenceville, 13—14. VIII. 51, 2 ind. (L); Grand Bank, 1. VIII. 51, 2 ind. (L); St. Pierre, 7—12. VIII. 51, 1 ind. (f. macr.) 11. VIII. 51, 1 ind. (L); Michelon, 8. VIII. 51, 2 ind. (L); Ramea Island, 15—26. VI. 49, 1 ind. (L). — *W.* Corner Brook, 14. VIII. 49, 2 ind. (L); Deer Lake, 17—19. VIII. 49, 1 ind. (L). — *N.W.* Port au Choix, 4—5. VIII. 49, 3 ind. (L); Eddies Cove West, 28—29. VII. 49, 1 ind. (L). — *N.E.* Hamden, 24. VII. 51, 1 ind. (L); Twillingate, 4—8. VII. 51, 3 ind. (L); Fogo Tilting, 29—30. VI. 51, 1 ind. (L). — *E.* Gambo, 25. VIII. 49, 1 ind. (L); Port Rexton, 30—31. VII. 51, 1 ind. (L); Shoal Harbour, 29. VII. 49, 1 ind. (L); Clarenville, 31. VII. 51, 1 ind. (L). — *C.* Millertown Junction, 22. VIII. 49, 2 ind. (L).

*E c o l o g y*: On grasses and herbaceous vegetation. From many localities in Newfoundland.

*D i s t r i b u t i o n*: Circumpolar holarctic species. With wide distribution in the Euro-siberian subregion. Only in the northern parts of the nearctic region. Recorded from Alaska, Canada, Greenland, Maine, Colorado.

*N. limbatus* Dahlb.

*Nabis limbatus* Dahlb. K. V. A. Handl. 1850, 227. — v. D. 279. — Bl. 598. — Mo. 16. — Strickl. 199.

*C.* Millertown Junction, 21. VIII. 49, 1 ind. (K).

*D i s t r i b u t i o n*: Newfoundland, Quebec, Ontario, Alberta. In the States from New England to Colorado. Probably an introduced palearctic species. In the palearctic region with nearly as wide a range as *N. flavomarginatus*.

MIRIDAE

*Pithanus* Fieb.

*P. maerkeli* (H. S.)

*Capsus maerkelii* H. S. Wanzenart. Ins. I, 4, 1839, 78. — *Pithanus maerkelii* v. D. 313. — Bl. 680.

*S.* Grand Bank, 2—3. VIII. 51, 1 ind. (L); Michelon, 9. VIII. 51, 1 ind.; Deer Lake, 25. VII. 51, 1 ind. (L); *E.* Clarenville, 31. VII. 51, 1 ind. (L).

*E c o l o g y*: Only f. brach. On sandy localities, on low dense vegetation, but also on humid meadowland with *Carex*, *Juncus*, etc.; on cultivated land.

*D i s t r i b u t i o n*: Palearctic species, introduced from Europe. Recorded from Newfoundland, Nova Scotia and some eastern States: Maine, New York.