

**E c o l o g y:** On swamps, marshes, and bogs. — In L'Asne au Loup by a pool with *Carex* and *Sphagnum*.

**D i s t r i b u t i o n:** This species has a remarkable geographical distribution. It was described from Labrador and later found in Quebec, Bradore Bay. It is also known from the fjelds of Scandinavia.

*T. paludum* J. Sahlb.

*Teratocoris paludum* J. Sahlb. Not. Soc. F. Fl. Fenn. 11, 1871, 291. — Bl. 678.

S.E. St. Johns, 13. VIII. 49, 2 ind. (L).

**E c o l o g y:** On a quagmire.

**D i s t r i b u t i o n:** Palearctic species from northern Europe and Siberia. Introduced to North America (Blatschley); known from New York, Illinois and Minnesota.

#### *Capsus* Fabr.

*C. ater* (L.)

*Cimex ater* L. Syst. Nat. ed. 10, 1758, 447. — *Capsus a.* v. D. 338, — Bl. 753. — Mo. 20.

W. South Branch, 3. VII. 49, 5 ind. (P).

**E c o l o g y:** On a moist meadow. Two specimens belong to var. *semiflavus* (L.).

**D i s t r i b u t i o n:** Palearctic species, eastward to western Siberia. Introduced from Europe. From Newfoundland and Quebec to Alberta; from New England west to South Dakota and south to Florida.

#### *Orthops* Fieb.

*O. scutellatus* Uhl.

*Orthops scutellatus* Uhl. Bull. U. S. Geol. Geogr. Surv. III, 1877, 420. — *Lygus campestris* v. D. 348. — Bl. 767. — Mo. 18. — *L. scutellatus* Strickl. 202. — Kelton, Can. Ent. LXXXVII, 540.

S. Burgeo, Grandy Brook, 23. VI. 49, 2 ind. (K). — W. South Branch, VII. 49, 3 ind. (K); Piccadilly, 7. VII. 49, 3 ind. (L); Piccadilly, Port au Peninsula, 7. VII. 49, 4 ind. (P); Spruce Brook, 8—9. VII. 49, 17 ind. (P); Lomond, Bonne Bay, 13. VII. 49, 5 ind. (P). — N.W. Williamsport, 13. VII. 49, 3 ind. (L).

**E c o l o g y:** On *Umbelliferae*, in Piccadilly found on *Heracleum*.

**D i s t r i b u t i o n:** Nearctic species with northern distribution. From Newfoundland and New England to Alaska and the Pacific.

#### *Lygus* Hahn

*L. (Lygus) pabulinus* L.

*Cimex pabulinus* L. Fauna Sv. 1761, 253. — *Lygus p.* v. D. 340. — Walley, Can. Ent. LVII, 76. — Mo. 18. — Strickl. 201.

N.W. Cow Head, 11. VIII. 49, 2 ind. (P); Doctors Brook, 31. VII. 49, 1 ind. (L). — E. Port Rexton, 30. VII. 51, 1 ind. (L). — C. Lake St. John, 27. VII. 51, 1 ind. (L).

**E c o l o g y:** In meadows along river banks, in shady places, in woods.

**D i s t r i b u t i o n:** Holarctic species. Northern and middle palearctic region. In the northern part of the nearctic region, from the Atlantic to the Pacific. In the States from New England west to Michigan and southwest to North Carolina.

*L. (Neolygus) communis* Kngt

*Lygus communis* Kngt, Can. Ent. XLVIII, 1916, 346. — v. D. 343. — *L. (Neolygus) c.* Bl. 780. — Walley, Can. Ent. LXII, 75. — Mo. 19. — Strickl. 202.