

(P); Steady Brook, Humber, 10. VII. 49, 7 ind. (P); Lomond, 13. VII. 49, 1 ind. (P). — N.W. Williamsport, 13. VII. 49, 1 ind. (L); — N.E. Springdale, 20—21. VI. 51, 1 ind. (L). — E. Gambo, 26. VIII. 49, 1 ind. (P). — C. Gander, 1. IX. 49, 1 ind. (L).

E c o l o g y: On a *Kalmia*-bog with open vegetation of grass and *Carices*, on swamps with *Carex-Juncus*, on shaded meadows in woodland.

D i s t r i b u t i o n: From Newfoundland across Canada to British Columbia. From some northern States.

L. rubroclarus Kngt

Lygus vanduzeei var. *rubroclarus* Kngt, Corn. Univ. Agri. Expt. Sta. Bull. XXXIV, 1917, 567. — Mo. 18. — *L. rubroclarus* Strickl. 202. — *Liocoris rubroclarus* Kelt. Can. Ent. LXXXVII, 1955, 552.

S.E. Portugal Cove, 6. VI. 49, 2 ind. (K). — S. Rencontre West, 17—18. VI. 49, 1 ind. (L); Hare Bay, 21. VI. 49, 1 ind. (L). — W. Steady Brook, 10. VII. 49, 1 ind. (L).

D i s t r i b u t i o n: From Newfoundland across Canada to Alberta. Northwestern States.

L. columbiensis Kngt

Lygus columbiensis Kngt, Corn. Univ. Agri. Expt. Sta. Bull. XXXIV, 1917, 567. — *Liocoris c.* Kelt. Can. Ent. LXXXVII, 1955, 554.

S. Rencontre West, 17—19. VI. 49, 1 ind. (L).

D i s t r i b u t i o n: Newfoundland; Manitoba, Saskatchewan, Alberta, British Columbia, Northwest Territories; Alaska.

Salignus Kelton

S. distinguendus (Reut.)

Lygus distinguendus Reut. Pet. Nouv. Ent. I, 1875, 541. — v. D. 349. — Strickl. 201. — *Salignus d.* Kelton, Can. Ent. LXXXVII, 536.

W. Deer Lake, 18. VIII. 49, 1 ind. (L).

E c o l o g y: On cultivated land.

D i s t r i b u t i o n: Nearctic species with wide range. Newfoundland; Alaska, Yukon, western states of Canada; California, Utah, Colorado, New Mexico. — Siberia.

Camptobrochis Fieb.

C. nebulosus Uhl.

Camptobrochis nebulosus Uhl. Haydens Surv. Terr. for 1871, 1872, 417. — v. D. 354. — Bl. 889. — Mo. 22.

S. Hare Bay, 21. VI. 49, 2 ind. (L).

D i s t r i b u t i o n: Nearctic species with wide range. South part of Newfoundland, Quebec, Ontario, New England west to the Pacific and south to Florida and Texas.

Monalocoris Dahlb.

M. americanus Wagn. & Slat.

Monalocoris filicis v. D. 361. — Bl. 874. — Walley, Can. Ent. LXII, N.B. 76. — Mo. 22. — *M. americanus* Wagn. and Slat. Proc. Ent. Soc. Wash. 54, 1952, 279. — Strickl. 200.

S. Rencontre West, 17—19. VI. 49, 1 ind. (L); Grand Bruit, 13. VI. 49, 1 ind. (L).

E c o l o g y: On *Dryopteris* and *Osmunda*.

D i s t r i b u t i o n: Nearctic species with wide range. From Newfoundland and Quebec to British Columbia. Generally distributed in the States.