

*Dicyphus* Fieb.*D. agilis* (Uhl.)*Itolocoris agilis* Uhl. Bull. U.S. Geol. Geogr. Surv. III, 1877, 425. — *Dicyphus a.* Bl. 909. — Mo. 22. — Strickl. 200.

S.E. Labrador: Pinware River, 19. VII. 51, 1 ind. (L).

N.W. Doctors Brook, 31. VIII. 49, 1 ind. (K)

E c o l o g y: In forest. The species lives on *Rubus*.

D i s t r i b u t i o n: Nearctic species, fairly widespread. From Newfoundland, Quebec and New England to the Pacific, southwest to North Carolina.

*Mecomma* Fieb.*M. gilvipes* (Stål)*Leptomerochoris gilvipes* Stål, Stett. Ent. Zeit. IX, 1858, 187. — *Mecomma g.* v. D. 398. — Bl. 852. — Walley, Can. Ent. LXII, 76. — Mo. 21.

S.E. Labrador, Grand Falls, 15. VII. 51, 1 ind. (L).

N.W. Doctors Hill, 31. VII. 49, 1 ind. (L).

E c o l o g y: On marshes and meadows on shores.

D i s t r i b u t i o n: Nearctic species with northern distribution. From Newfoundland, Quebec and New England west to the Pacific.

*Cyrtorrhinus* Fieb.*C. caricis vagus* Knigt*Capsus caricis* Fall. Cimic. Suec, 1807, 102. — *Cyrtorrhinus c.* v. D. 824. — Bl. 853. — *C. c. vagus* Knigt, Hem. Connecticut, Miridae, 1923, 511.

S.E. St. Johns, 13. VIII. 49, 3 ind. (L); Come-by-Chance, 24. VIII. 49, 1 ind. (L). — W. Deer Lake, 18. VIII. 49, 2 ind. (L). — E. Goobies, 15. VIII. 51, 1 ind. (L). — C. Grand Falls, 26. VII. 51, 1 ind. (L).

E c o l o g y: On bogs and marshes, on shores of lakes; on thick vegetation of *Carex* and *Equisetum*.D i s t r i b u t i o n: Probably a holarctic species. The subspecies is recorded from Massachusetts, New York, New Jersey and Virginia, the typical *caricis* from Colorado.*C. n. sp.*

E. Port Rexton, 31. VII. 51, 2 Exx. (L). — C. Grand Falls, 26. VII. 51, 1 ind.

E c o l o g y: On high thick vegetation of *Carex*, and *Equisetum* in *Sphagnum* pools.*Plagiognathus* Fieb.*P. obscurus* Uhler var. *fraternus* Uhl.*Plagiognathus obscurus* Uhler, Haydens Surv. Terr. for 1871, 1872, 418. — *P. fraternus* Uhler, Hemipt. Colorado, 1895, 51. — *P. fraternus* v. D. 411. — Bl. 931. — Walley, Can. Ent. LXII, 76. — Mo. 23. — Strickl. 200.

C. Millertown Junction, 21. VIII. 49, 1 ind. (K).

D i s t r i b u t i o n: Nearctic species with wide range. Newfoundland, Nova Scotia, Quebec, Ontario, Alberta, British Columbia. From New England west to Kansas, Colorado.

*P. politus* Uhl.*Plagiognathus politus* Uhl. Hemipt. Colorado, 1895, 52. — v. D. 411. — Bl. 932. — Walley, Can. Ent. LXII, 76. — Mo. 24.