

As *L. pabulinus* (L.), but robuster, dissimilarly coloured and provided with dissimilar male genitalia. Easily recognized by the golden yellow colouring and the sanguineous clypeus.

L. honshuensis n.sp.

Length 5.25–6.25 mm. Externally much as *L. contaminatus* (FN.). Yellowish green. Apex of clypeus slightly darkened. Apex of 2nd antennal joint and 3rd and 4th joints slightly infusate. A transverse dark fuscous band along inner part of apical margin of corium; membrane smoky with some darker shadows, basal inner angle dark fuscous.

Body as in *L. contaminatus*. Crown $0.87 \times$ (σ) or $1.0 \times$ (φ) as broad as eye, with a longitudinal furrow, hind margin distinctly carinate. 2nd antennal joint $1.0 \times$ (σ) or $0.98 \times$ (φ) as long as basal width of pronotum. Pronotum densely and finely punctate. Elytra densely punctate, hair covering yellowish, relatively long. Rostrum extending to hind coxae. Male genitalia: Left stylus (fig. 2 a) slender, nearly semicircularly curved in outline. Right stylus (fig. 2 b) relatively short and stout, hypophysis claw-like, sensory lobe distinctly produced. Chitinized aedeagal appendage long and slender, strongly curved apically (fig. 2 c).




Fig. 2. *Lygus honshuensis* n.sp.: a left stylus; b right stylus; c chitinized aedeagal appendage. – *L. pulchellus* Rr.: d–f same. – *L. disponsi* n.sp.: g left stylus; h hypophysis of same, dorsal aspect. – Orig.