

gracile, the antennae are much longer and the hair covering longer. *P. sinuaticollis* Rt. has the lateral margins of the pronotum strongly insinuated and the antennae longer. *P. setulosus* Hv. is provided with long erect hairs on the upper surface, the vertex is twice as broad as the eye and the cuneus lacks the silvery band.

*P. niger* POPP.

Kyushu, Wakasugi-yama, near Fukuoka, 3 spec., 13. X. 1957, S. MIYAMOTO.  
– Endemic.

*Cyrtorhinus lividipennis* Rt.

Kyushu, Otomari (Osumi), 2 spec., 30. VIII. 1951, S. MIYAMOTO. – Oriental.

*Orihotylus flavosparsus* (C.SB.)

Kyushu, Fukuoka, 3 spec., 19. VI. 1958. S. MIYAMOTO. – Holartic; a new record for Japan.

*Halticus micantulus* Hv.

Kyushu, Korasen, 3 spec., 9. VI. 1957, S. MIYAMOTO; Kyushu, Omura, Hizen, 1 spec., 4. VIII. 1952, S. MIYAMOTO. – Endemic.

*Psallus wagneri* Oss.

Hokkaido, Ikeda, 2 spec., 2. VII. 1958, S. MIYAMOTO. – Euro-Siberian.  
Previously known from Sweden, Finland and West Siberia.

*P. senjoensis* n.sp.

Length 4 – 4.5 mm. Head, pronotum and scutellum dark brown with a faint greyish tinge, relatively shining. Antennae dark brown. Elytra subopaque, greyish; commissural margin of clavus and a longitudinal dash on corium faintly fuscous; membrane smoky, veins light grey. Under surface and legs greyish brown, partly tinged with greenish. Femora minutely spotted with fuscous; tibial spines black, arising from distinct black spots.

Body 3.3 × (♂) or 2.6 × (♀) as long as broad. Vertex 1.83 × (♂) or 2.63 × (♀) as broad as eye. Eyes relatively small. Antennae long and gracile, proportions between joints 9 + 40 + 30 + ? (♂) or 9 + 29 + ? + ? (♀), 2nd joint 1.14 × (♂) or 0.83 × (♀) as long as basal width of pronotum. Pronotum 1.5 – 1.6 × as broad basally as head, 2.33 × as broad as long. Elytra extending well beyond abdomen. Hair covering of upper surface yellowish brown, obliquely erect and dense; in addition shorter smooth whitish hairs present on elytra. Rostrum extending to hind coxae. Claws distinctly curved about as in *P. luridus* Rt. Male genitalia: Left stylus (fig. 4 c) with hypophysis rather straight; sensory lobe with a short