

New or lesser known species of the genus *Pilophorus* Hhn. (Het., Miridae)

R. LINNAVUORI

1. *Pilophorus setulosus* Hv.

♂ Length 3.5–4 mm. Head blackish brown, tinged with rufous in lower part. Pronotum and scutellum blackish, the latter with 3 groups of silvery hairs. 1st and 2nd antennal joints light reddish brown, the apical third of the latter black, other joints whitish. Elytra dull cinnamon, with apex of clavus and of exocorium shining; with 2 transverse bands of silvery hairs, the posterior band straight; also base of cuneus with silvery hairs; membrane brownish smoky, becoming lighter apically. Under surface blackish; abdomen with an oblique band of silvery hairs on either side. Legs reddish brown, coxae apically whitish.




Fig. 1. *Pilophorus setulosus* Hv.: a head and pronotum. — *P. erraticus* n. sp.: b same. — *P. sinuaticollis* Rt.: c same. — *P. ponderosus* n. sp.: d same. — Orig.

Fig. 1 a. Body elongate, $3.6 \times$ as long as broad. Crown $1.86 \times$ as broad as eye. Proportions between antennal joints $9 + 40 + 17 + 13$ (1 unit = 0.033 mm.); 2nd joint $1.14 \times$ as long as basal width of pronotum, relatively thin and only slightly thickening apically; 3rd joint $0.43 \times$ as long as 2nd. Pronotum $1.67 \times$ as broad as long, strongly tapering anteriorly, lateral margins moderately