



Fig. 34. *Orthotylus mollis* sp. n.: a pygophore from side; b - c right and d left stylus; e penis. - *O. repandus* sp. n.: f right stylus; g processes of pygophore; h left stylus; i processes of vesica. - *O. compactus* sp. n.: j pygophore from side; k right and l left stylus; m processes of vesica; n ventral process of same (broken in Fig. m).

✓ *O. compactus* sp. n.

Length 3 - 4.5 mm. Rather shiny, greenish yellow. Antennae and legs yellow-brown. Elytra green, membrane pale, veins green.

Remarkably robust, elongately ovate, 2.7 - 3.0 × as long as broad at pronotum. Hair covering pale. Head 0.7 × as broad as pronotum, vertex basally sharply marginate and provided with a median longitudinal depression, ocular index 1.5 (♂) or 2.18 (♀). Proportions between antennal joints. 7 : 30 : ? : ?, 2nd joint as long as basal width of pronotum. Rostrum extending to near hind coxae. Pronotum slightly more than twice as broad as long, calli distinctly elevated. Male genitalia: right side of genital segment (Fig. 34 j) with numerous black teeth; right stylus (Fig. 34 k) small, ovate; left stylus as in Fig. 34 l, sensory lobe completely rounded; vesical appendages as in Fig. 34 m - n.

Equatoria: Torit - Kapoeta, 1 ♂, type and 1 paratype, 26. III. 1963; 52, 1 paratype. Not closely related to the preceding species and possibly representing another group.

3. Subgenus *Melanotrichus* Rt. (= *Halocapsus* Pt.)

✓ *O. (M.) martini* Pt. - 14, 1 ex. At lamp. Eremian, previously known from Algeria.

✓ *O. (M.) pusillus* Rt. - 14, several exx. On *Suaeda monoica* bushes in salt marshes. Eremian (Israel, Egypt, Arabia, Eritrea, Somalia).

✓ *O. (M.) omanensis* sp. n.

Length 3.2 mm. Rather shiny. Green. Antennae yellowish. Legs yellowish green.

Robust, parallel-sided, 2.5 × as long as broad. Hair covering pale, also silvery adpressed pubescence present. Head about 0.7 × as broad as pronotum, in apical view 1.4 × as broad as high, base of vertex distinctly marginate, ocular index 2.0. Antennae long, proportions between joints 6 : 27 : 22 : ?, 2nd joint 0.9 × as long as basal width of pronotum. Rostrum extending to hind coxae. Male genitalia: right stylus (Fig. 35 c - d) incrassate, with a long hypophysis; left stylus (Fig. 35 a - b) with some minute teeth in sensory lobe, hypophysis long; penis as in Fig. 35 e.

Oman, 1 ♂, type, in my collection.

Near *O. hirtulus* Wgn., but smaller, without black hairs, sensory lobe of left stylus with a small apical spine and right stylus dissimilar. *O. arabicus* Wgn. is smaller and more elongate, the base of the vertex is not marginate and the styli are differently shaped.

✓ *O. (Melanotrichus) monticolus* sp. n.

Length 3.5 mm. Relatively opaque. Pale yellowish. Antennae and legs yellowish. Membrane smoky, veins yellowish.

Robust, 2.6 × as long as broad. Semi-erect hair covering completely pale, smooth silvery hairs sparse. Head nearly as broad as pronotum, in apical view 1.5 × as broad as high, vertex distinctly marginate, ocular index 1.50 - 1.63 (♂). Proportions between antennal joints 5 : 22 : 18 : 12, 2nd joint slightly longer than basal width of pronotum. Rostrum extending to hind coxae. Pronotum 1.8 × as broad as long. Hind tibia 3.4 × as long as tarsus. Male genitalia much as in *O. halabicus*, but hypophysis of left stylus (Fig. 35 f - g) thinner and penis (Fig. 35 i - j) robust; right stylus (Fig. 35 h) straighter.