



Fig. 35. *Orthotylus omanensis* sp. n.: a - b left and c - d right stylus; e penis. - *O. monticolus* sp. n.: f left stylus; g same from above; h right stylus; i penis from side; j same, ventral aspect. - *O. halaibicus* sp. n.: k left stylus; l same from above; m sensory lobe of same; n right stylus; o penis from side; p same, ventral aspect. - *O. halaibicus abbreviatus* ssp. n.: q - r left and s - t right stylus. - *O. ericinellae* Pop.: u right and v left stylus; x penis.

Kassala: Erkowit, 1 ♂, type and 1 ♂, paratype, 5 - 10. VII. 1961. At lamp.

Recognized by the pale hair covering, the broad head with large eyes, etc.

✓ *O. (Melanotrichus) halaibicus* sp. n.

Length 3.5 mm. Shiny. Yellowish or yellowish green. Membrane slightly smoky, veins greenish. Antennae and legs yellowish.

Body robust, only 2.6 × as long as broad at pronotum. Hair covering remarkably long, mainly pale, on cuneus darker, silvery adpressed hairs sparse. Head broad, 0.7 × as broad as pronotum, in apical view 1.40 - 1.57 × as broad as high, vertex basally distinctly marginate, eyes prominent, ocular index 1.40 - 1.67 (♂) or 2.2 (♀). Proportions between antennal joints 6 : 27 : 21 : 11, 2nd joint slightly shorter than basal width of pronotum. Rostrum extending to hind coxae. Pronotum 2.4 × as broad as long. Elytra longer than abdomen. Hind tibia 3.6 × as long as tarsus. Male genitalia: right stylus as in Fig. 35 n; left stylus (Fig. 35 k - m) with a rather thick hypophysis, sensory lobe with a few minute apical teeth. Penis (Fig. 35 o - p) remarkably gracile.

10, several paratypes; 11, several paratypes; 12, 1 paratype; Suakin, 1 ♂, type and 1 paratype, 5. XII. 1962. On *Suaeda monoica* in salt marshes on the Red Sea coast.

Recognized by the robust body, the long hair covering, the shape of the head etc.

✓ *O. (Melanotrichus) halaibicus* Lv. ssp. *abbreviatus* ssp. n.
Length 3 mm. Much smaller and more gracile than the

nominate form, ocular index 1.6 - 1.7 (♂), 2nd antennal joint slightly longer than basal width of pronotum. Genitalia (Fig. 35 q - t) as in the nominate form, but the expanded apical part of right stylus smaller.

Arabia, near Lith, 1 ♂, type (British Museum) and 1 ♂ paratype (in my collection), 1945, Uvarov.

4. Subgenus *Litocoris* Fb.

✓ *O. (Litocoris) ericinellae* Pop.

For description see Poppius 1914: 69. Male genitalia as in Fig. 35 u - x. *O. ericinellae* differs in the simple structure of the penis from the Palaearctic species of the subgenus.

Material studied: Ethiopia, 98, 3 exx. On *Erica arborea*. Otherwise known from Kilimandjaro on *Ericinella manni*.

Pseudoloxops Kk.

✓ *P. sudanicus* sp. n.

Length 4 mm. Like *P. coccineus* (M. - D.), but smaller, with less intense red markings, with pale hind femora marked apically only with a narrow red ring and with dissimilar genitalia (Fig. 36 a - f).

Measurements: ocular index 1.33 (♂) or 2.0 (♀). Proportions between antennal joints 11 : 41 : 14 : 10, 2nd joint 1.2 - 1.3 × as long as basal width of pronotum. Rostrum extending to hind coxae.

Kassala: Kassala, 1 ♂, type, 30. XI. 1962; 72, 2 paratypes. At lamp.