



Fig. 52. *Plagiorrhamma concolor* (Rt.): a - b left stylus; c theca. - *P. ruficollis* sp. n.: d - f same. - *P. monticolus* sp. n.: g right and h - i left stylus; j theca; k apex of vesica. - *P. quadripunctatus* sp. n.: l apex of elytron. - *P. albofasciatus* (Motsch.): m left stylus; n theca; o apex of vesica. - *P. curtipes* sp. n.: p head and thorax from side; q - r left stylus; s theca; t apex of vesica.

- 29 (30) Antennae remarkably thicker .. *P. discoidalis*  
 30 (29) Antennae gracile (as, e.g., in *costai*) ..... 31  
 31 (32) Theca with a single, serrate appendage (Fig. 53 h - j). ..... *P. similis*  
 32 (31) Theca with a bifurcate appendage (Fig. 53 f - g) ..... *P. jocosus*

*P. poseidon* (Kk.), unknown to me, is excluded from the key.

✓ *P. concolor* Rt.

Theca (Fig. 52 c) distinctly curvate. Left stylus as in Fig. 52 a - b.

50 km. E of 9, 1 ex.; near 43, 1 ex.; 96 - 40, 1 ex.; 35, 2 exx.; 35 - 36, 1 ex.; 40, 2 exx.; 32, several exx.; 33 - 34, several exx.; 45, several exx. At lamp. Apparently Eremian. Previously known from the Pontomediterranean area.

✓ *P. ruficollis* sp. n.

Like *P. concolor*, but 1) head, pronotum and scutellum more intensely reddish brown or ferrugineous, also antennae and legs more intensely reddish, 2) eyes larger, ocular index 1.4 - 1.7 (♂) (in *concolor* 1.73 - 2.0 (♂) or 2.0 (♀)), 3) 2nd antennal joint about 1.7 × as long as diatone (in *concolor* 2.0 - 2.53 ×), 4) theca (Fig. 52 f) thicker and straighter and 5) hypophysis of left stylus (Fig. 52 d - e) dissimilar, claw-like.

6, 1 ♂ paratype; Kassala: Kassala - Haiya, 1 ♂, type and 1 ♂ paratype, 1 - 3. XII. 1962; 24 - 24a, 1 ♂ paratype. At lamp.

✓ *P. punctatulus* sp. n.

Fig. 51. Length 3.5 - 4 mm. Shiny. Head dark reddish brown, subopaque. Antennae orangish, 4th joint dark. Pronotum and scutellum black. Elytra dark brown, apex of corium with a squarish or roundish whitish spot rarely also base of corium slightly paler, uncus dark purplish, membrane uniformly dark brown. Under surface dark reddish brown. Legs reddish brown, middle and hind coxae pale.

Robust, body 3.3 × as long as broad. Hair covering golden, adpressed. Head strongly shagreened, without a median sulcus, tylus strongly prominent, ocular index 1.4 (♂) or 2.27 (♀). Antennae relatively thick, proportions between joints 9 : 30 : 24 : 14, 1st joint 0.53 × as long as diatone, 2nd slightly longer than basal width of pronotum. Pronotum strongly broadening caudad, nearly twice as broad as long, anterior part shagreened, disk only very faintly shagreened, but finely punctate and rugose. Scutellum transversely wrinkled. Elytra finely punctate, longer (♂ ♀) than abdomen; hair covering of upper surface most distinct on elytra.

63 - 62, 1 paratype; 68, 1 paratype; Lotti forest, 1 ♂, type, 14 - 17. III. 1963; 66 - 64, 3 paratypes. At lamp. Also known from Zaire (Katanga).

✓ *P. monticolus* sp. n.

Length 3.25 - 4 mm. Opaque or subopaque. Pale reddish brown. Clavus and inner apical angle of corium and some-