



Zanchisme illustris Reuter. 1. Head of male, frontal view; 2. Head of female, frontal view; 3. Head and pronotum, female, lateral view; 4. Pronotum, male, lateral view; 5. Pronotum, female nymph, lateral view; 6. Right clasper, ventral view; 7. Right clasper, outer lateral view; 8. Left clasper, dorsal view; 9. Left clasper, outer lateral view; 10. Tip of aedeagus, outer lateral view; 11. Tip of aedeagus, dorsal view.

Frons inconspicuously striated above clypeus, vertex smooth; eyes relatively large in males, smaller in females, reaching posterior margin of head. Antenna with second segment longest, first shortest, segments of subequal thickness and somewhat subfusiform; base of antenna partially surrounded by eyes in males, about midway from eyes to clypeus in females. Rostrum slender, thinner than antenna; apex reaching to before middle coxa; first segment partially hidden by bucculae. Hemielytra opaque, costal margin concave; embolium widening to apex; cuneus longer than wide, inner margin semicircular; with sparse long semi-decumbent pubescence and silvery scale-like hairs arranged in bands and stripes on clavus and corium; membrane rounded apically, two-celled. Mesoscutum exposed; scutellum triangular, with or without silvery scale-like hairs. Coxae and trochanters well developed; mid- and hindfemora slightly curved, forefemora straight. Abdomen constricted at base, gradually widening from base to last or genital segment.

Type of genus *Schizonotus dromedarius* Reuter 1892.

***Zanchisme illustris* Reuter.**

Zanchisme illustris Reuter. Ofv. F. Vet. Soc. Forh. 49 (5):11.