

*Paratypes*: Same data as for holotype, 3 ♀; Carbondale, Illinois, May 18, 1955, S. C. Chandler, on Wisteria vine, 1 ♂ (teneral). All types deposited with the Illinois Natural History Survey. This species is named for the collector, Mr. Stewart C. Chandler.

### **Lopidea wisteriae** new species

As with the above species, closely related to *L. amorphae* and *L. reuteri*, but separated by means of male genital characters.

*Male*.—Size subequal to that of *L. reuteri*; coloration as in *L. chandleri* with the following exceptions: Head with a broad black stripe at each side of the front between antennae; pronotum with infuscated broad apical transverse band and fuscous inverted "V" above scutellum; scutellum darker; each hemelytron with clavus and inner half of corium generally infuscated; femora more reddish; abdomen with ninth segment somewhat infuscated; genital structures distinctive as illustrated.

*Female*.—Slightly larger and more robust than male; coloration similar to male; venter of abdomen red with black around ovipositor on segments 6 and 7, segment 8 black near ovipositor and on apical margin, 9 black on basal and apical margins with disc reddish, third valvulae black.

*Holotype*: ♂, Carbondale, ILLINOIS, July 7, 1955, H. H. Ross and T. E. Moore, on Wisteria vine. Deposited at the Illinois Natural History Survey.

*Paratypes*: Same data as for holotype, 19 ♂, 12 ♀. Specimens deposited with the Illinois Natural History Survey, the U. S. National Museum, and the British Museum (Natural History).

#### REFERENCES CITED

- KNIGHT, H. H. 1918. Ent. News, 29: 172-176.  
—. 1941. Bull. Ill. Nat. Hist. Surv., 22: 84-91.