

Holotype: Male, MEXICO: OAXACA, 10 mi. n. Miltepec, July 26, 1974, Clark, Murray, Ashe, Schaffner. Deposited in the National Museum of Natural History, Washington, D.C. *Allotype*: Female, same data as holotype. Deposited in the National Museum of Natural History. *Paratypes*: 29 males, 16 females, same data as holotype: 6 males, 6 females, MEXICO: PUEBLA, 4.4 mi. sw. Acatepec, July 26, 1974. Clark, Murray, Ashe, Schaffner; male, 9 females, same locality as holotype except taken August 4, 1976, Peigler, Gruetzmacher, R&M Murray, Schaffner. Deposited in the collection of the Department of Entomology, Texas A&M University, College Station, Texas; in the Museo de Historia Natural de la Ciudad de Mexico, Mexico; in the J.C.M. Carvalho collection, Rio de Janeiro, Brazil; and in the Canadian National Collection, Ottawa, Ontario.

All male specimens collected near Miltepec had the tergal process of the genital capsule spinelike, whereas those specimens taken near Acatepec possessed a bifurcate tergal process. No other significant differences in the genitalia could be discerned.

This species is named in honor of Pedro Reyes-Castillo of the Instituto de Ecologia, Museo de Historia Natural de la Ciudad de Mexico, who has been of invaluable assistance concerning my interest in the Miridae of Mexico.

***Fulgenticapsus acatepeci* n. sp.**

(Fig. 3)

Characterized by the light brown to brown scutellum; also, in the case of the male, by the bifurcate tergal process and by the shape of the right paramere.

Male (measurements taken from 4 specimens, those of holotype given first followed in parenthesis by average and ranges): Length, 3.80 mm (3.86 mm, 3.80 – 3.92 mm); width, 1.54 mm (1.56 mm, 1.52 – 1.60 mm). Head length 0.14 mm (0.15 mm, 0.14 – 0.16 mm); width through eyes, 0.80 mm (0.81 mm, 0.80 – 0.82 mm); width, 0.36 mm (0.35 mm, 0.34 – 0.36 mm). Length of antennal segment I, 0.26 mm (0.25 mm, 0.22 – 0.26 mm); II, 1.10 mm (1.08 mm, 1.04 – 1.10 mm); III, 0.84 mm (0.88 mm [one specimen]); IV, 0.38 mm (0.34 mm [one specimen]). Pronotal length, 0.66 mm (all specimens); width at base, 1.30 mm (1.31 mm, 1.30 – 1.34 mm). Cuneal length, 0.64 mm (0.68 mm, 0.61 – 0.70 mm); width at base, 0.52 mm (0.54 mm, 0.50 – 0.56 mm).

General coloration light brown above, dark fuscous to black beneath; head light brown with reddish areas on apex of jugum and sometimes on