

shoek Pass Summit, 5400 ft. elevation, 25 mi. S. Harrismith, 4 Mar. 1968, T. Schuh, J. A. & S. Slater, M. Sweet (SANC).

PARATYPES: *Natal*—1 macropterous ♂, Cathedral Peak, Jan. 1964 (Capener); 1 macropterous ♂ (?), Drakensberg, 1-22.I.1927 (Turner); 1 macropterous ♂, same data as holotype. *Transvaal*—1 macropterous ♀, Wylies Poort, 10.2.41 (Capener) (SANC, BM [NH], RTS).

This species is named for its occurrence in Natal.

Zanchiella natalensis is most closely related to *Z. capensis* and *Z. bowkeriae*. It can be recognized by the pronotum being light on the anterior third and dark on the posterior two-thirds, in combination with the dark transverse macula on the posterior half of the corium.

***Zanchiella sweeti*, new species**

Figure 18

MACROPTEROUS MALE: Elongate, nearly parallel sided; head (excepting posterior margin of vertex and genae), pronotum, scutellum, clavus along posterior two-thirds of commissure, and transverse macula on corium at level of base of membrane orangish brown; genae, posterior half of vertex, hemelytra (excluding membrane), thoracic venter, legs, and labium very light green; abdominal venter green; proximal fifth of antennal segment 1 light green, distal four-fifths red; antennal segment 2 brown on proximal end, remainder with broad reddish brown bands alternating with light coloration; antennal segments 3 and 4 brown; membrane light smoky gray.

Head polished, smooth, and shining; pronotum and scutellum obscurely transversely rugulose, weakly shining; hemelytra subhyaline, dull; membrane rugulose; dorsum with semierect, moderately long, light hairs; anterolateral angles of pronotum with single, long, erect, fine seta; antennae with short, decumbent, light vestiture; abdomen with moderately long, reclining, light hairs.

Frons weakly convexly produced between eyes; anterior margins of eyes weakly emarginate; labium just surpassing metacoxae; calli rather widely separated, demarcated posteriorly by weak furrow; posterior pronotal margin very slightly convex; punctures on clavus faint; tibiae with scattered, very fine, short, reclining light spines, hardly distinguishable from light tibial hairs; metatarsal segment 1 about half length of segment 2; segments 2 and 3 subequal in length.

MEASUREMENTS: Total length 3.24, maximum width .98, length