

- * *reuteri* Poppius (*Tyraquellus*), see *albofasciatus* (Motschulsky) **New Synonymy.**
similis Poppius (*Plagiorhamma*), 1914a, p. 55. East Africa; South Africa: Transvaal.
 * *tenuis* Odhiambo (*Hallodapus*), see *albofasciatus* (Motschulsky) **New Synonymy.**
transvaalensis, new species. South Africa: Natal, Transvaal.
verticicus Odhiambo (*Hallodapus*), 1967, pp. 1671–1673. Central African Republic.
vittatus Odhiambo (*Trichophthalmocapsus*), 1959c, pp. 661, 663–664, 685. **New Combination.** Uganda; Abyssinia.

KEYS TO SOUTH AFRICAN SPECIES OF *Hallodapus*

Macropterous specimens

1. Exocorium entirely cream or white 2
 Exocorium not entirely cream or white 3
2. Metafemora entirely cream or white *similis* (Fig. 30)
 Metafemora cream or white on proximal third, castaneous on distal two-thirds *pseudosimilis* (Fig. 28)
3. Hemelytra without complete white transverse fascia
 *quadrimaculatus* (Fig. 29)
 Hemelytra with complete white transverse fascia, femora not entirely castaneous 4
4. White transverse fascia of nearly uniform width
 *albofasciatus* (Fig. 27)
 White transverse fascia wide on corium, narrow on clavus
 *transvaalensis* (Fig. 31)

Brachypterous specimens

1. Posterior margins of hemelytra almost entirely white
 *transvaalensis* (Fig. 32)
 Posterior margins of hemelytra almost entirely castaneous
 *albofasciatus*

***Hallodapus albofasciatus* (Motschulsky)**

Figures 27, 163–165

Leptomerocoris albofasciatus Motschulsky, 1863, p. 86.

Tyraquellus reuteri Poppius, 1914a, pp. 51–52. **New Synonymy.**

Hallodapus tenuis Odhiambo, 1959c, pp. 664–668, 685. **New Synonymy.**

Trichophthalmocapsus chariensis Odhiambo, 1967, pp. 1678–1681. **New Synonymy.**

Hallodapus albofasciatus in South Africa can be recognized by the long, light colored, nearly erect pubescence on the dorsum, the