

## SUBFAMILY ISOMETOPINAE

## Tribe Diphlebiini

*Diphleps unica* Bergroth

Distribution. — Greenbrier

Seasonal Range. — 16 July

WV Host. — *Acer pseudoplatanus*

## Tribe Isometopini

*Corticoris pulchellus* (Heidemann)

Distribution. — Preston

Seasonal Range. — 8 Aug.

Remarks. — The only known West Virginia record is Aurora, Preston Co., the one Heidemann (1908) used in his original description of this species.

*Corticoris signatus* (Heidemann)

Distribution. — Mineral, Wood

Seasonal Range. — 10 June to 14

July

WV Host. — *Malus* sp. (crabapple)*Myiomma cixiiforme* (Uhler)

Distribution. — Grant, Mineral

Seasonal Range. — 10 June to 10

July

WV Host. — *Quercus coccinea*

## SUBFAMILY PHYLINAE

## Tribe Phylini

*Amblytylus nasutus* (Kirschbaum)

Distribution. — Berkeley, Grant,

Jefferson, Mineral, Monongalia

Seasonal Range. — 30 May to 10

June

WV Host. — grasses

*Atractotomus magnicornis* (Fallén)

Distribution. — Berkeley, Harrison,

Jefferson, Randolph

Seasonal Range. — 27 May to 3 July

WV Host. — *Picea abies**Campylomma verbasci* (Meyer)

Distribution. — Berkeley

Seasonal Range. — 25 Aug.

WV Host. — *Verbascum thapsus**Chlamydatus associatus* (Uhler)

Distribution. — Grant

Seasonal Range. — 4 June

*Criocoris saliens* (Reuter)

Distribution. — Berkeley, Jefferson,

Mineral, Preston

Seasonal Range. — 30 May to 25

June

WV Hosts. — *Galium aparine*, *G. mollugo*

*Keltonia sulphurea* (Reuter)

Distribution. — Wood

Seasonal Range. — 22 Aug.

*Lepidopsallus claricornis* Knight

Distribution. — Jefferson

Seasonal Range. — 15-16 May

WV Host. — *Pinus virginiana**Lepidopsallus miniatus* Knight

Distribution. — Jefferson

Seasonal Range. — 17 May

WV Host. — *Quercus alba**Lepidopsallus rostratus* Knight

Distribution. — Berkeley, Grant,

Hampshire

Seasonal Range. — 31 May to 4

July

WV Hosts. — *Pinus mugo*, *P. virginiana*

*Lepidopsallus rubidus* (Uhler)

Distribution. — Tucker

Seasonal Range. — 12 Aug.

WV Host. — *Salix* sp.