

Stenodema vicina (Provancher)

Distribution. — Greenbrier, Monongalia, Morgan, Pendleton, Pocahontas, Preston, Randolph, Tucker

Seasonal Range. — 8 Apr. to 24 Sept.

WV Hosts. — grasses

Trigonotylus coelestialium (Kirkaldy)

Distribution. — 13 counties; probably occurs statewide

Seasonal Range. — 29 May to 17 Aug.

WV Hosts. — *Poa pratensis*, other grasses

Tribe Herdoniini

Paraxenetus guttulatus (Uhler)

Distribution. — Greenbrier, Jefferson, Monongalia

Seasonal Range. — 16 July to 23 Aug.

WV Hosts. — *Acer pseudoplatanus*, *Staphylea trifolia*, *Viburnum recognitum*

SUBFAMILY BRYOCORINAE

Tribe Ecritotarsini

Monalocoris americanus Wagner & Slater

Distribution. — Grant, Jefferson, Morgan, Preston, Tucker

Seasonal Range. — 26 May to 20 Aug.

WV Host. — *Dennstaedtia punctilobula*

Sixeonotus tenebrosus (Distant)

Distribution. — Berkeley

Seasonal Range. — 2 Sept.

WV Host. — *Nasturtium officinale*

SUBFAMILY DICYPHINAE

Dicyphus agilis (Uhler)

Distribution. — Preston, Randolph, Tucker

Seasonal Range. — 26 June to 9 Aug.

WV Host. — *Rubus odoratus*

Remarks. — This is the species often referred to as *D. cucurbitaceus* (Spinola), which is Neotropical in distribution (Kelton 1980b).

Dicyphus famelicus (Uhler)

Distribution. — Pendleton, Tucker

Seasonal Range. — 9-26 June

WV Host. — *Rubus odoratus*

Dicyphus rhododendri Dolling

Distribution. — Greenbrier, Pendleton, Tucker

Seasonal Range. — 26 June to 15 July

WV Hosts. — *Rhododendron maximum*, *R. periclymenoides*

Macrolophus tenuicornis Blatchley

Distribution. — Pocahontas, Preston

Seasonal Range. — 24 June to 23 Aug.

WV Host. — *Dennstaedtia punctilobula*