

COMMENTS AND DESCRIPTIONS.

The insects mentioned below have all been communicated by Prof. C. F. Baker.

*Cylapomorpha gracilicornis* Popp.

The strongly thickened first antennal joint is described by Poppius as « aussen gerade, innen sehr stark erweitert », but in the single type from Los Baños the antennae had turned a little round their axis. In the normal position of the antennae the first joint is, seen from above, pear-shaped, very much thicker near the base than at the apex, and somewhat variable in size; it is also more or less distinctly pedunculate, the basal stalk being sometimes very short, scarcely perceptible, but in other specimens quite conspicuous. The other antennal joints are very long and more slender than in any other insect known to me, the last two joints being not much thicker than a cobweb's thread. All veins of the corium and clavus are distinct, the outer cell of the membrane is about half the length of the inner, and the vein separating the two cells of the membrane is continued beyond the cells. The cuneus is more or less narrowly yellow at the base.

*Trichofulvius zonatus* n. sp.

Piceo-niger, fascia lata leviter obliqua antemediana corii per trientem mediam clavi contindata, macula majuscula subquadrata ad apicem exteriorem corii ante basin cunei striolaque brevi longitudinali ad angulum apicalem interiorem corii albis, membrana infumata, iridescente; antennae dilute flavae, articulo primo fusco-nigro (duo ultimi desunt); rostrum fusco-testaceum; pedes flavo-albidi, coxis, trochanteribus ac femoribus fusco-nigris; caput et pronotum parce erecte concoloriter pilosa, scutellum et corium cum clavo densius semierecte longiuscule albido-pilosa. Caput pronoto medio paullo brevius, vertice oculo (♀) triplo latiore, prope basin utrinque mox intra oculos impressione brevi transversa praedito, articulo primo antennarum spatio interoculari (♀) paullo breviora, apicem clypei longius superante, secundo primo circiter triplo longiore et latitudine basali pronoti fere tertia parte longiore, rostro coxas posticas attingente, articulo primo marginem anticum prosterni paullum superante, secundo primo breviora, tertio primo subaeque longo, quarto tertio paullo breviora. Pronotum lateribus late levissime, basi late distinctissime sinuatum, callis confluentibus mox pone medium impressione transversa breviuscula com-