

antica præcipue in dimidio basali nonnihil incrassata; tibiae antice apice quam basi paullo latiores, posteriores apicem versus leviter angustatæ. Long. ♀ 3,5 mm.

Luzon : Los Banos.

Easily distinguished from *F. punctatus* Popp. by smaller size, different colour-markings, still larger pronotal calli, much more densely punctured corium, and by the cuneus which is completely fused with the corium, while the fracture is faintly indicated in *punctatus*.

Peritropis Poppiana BERGR.

In the description some words were left out by the printer. On page 119, line 8 and 9, the description should read : epipleura of corium fawn-colored, its exterior margin blackish, finely dotted with ochraceous; membrane fuscous, rather densely dotted with whitish.

This seem to be the commonest of the Philippine species of *Peritropis*. I have now received it from different localities in Luzon and from Mindanao (Butuan and Zamboanga). The head is as long as the pronotum, its antecular part seen from above distinctly longer than the eyes. The interior cell of the membrane is rounded or obtuse at apex, exterior cell as long as broad. The hind femora are sometimes entirely pale, excepting a dark subapical annulation. The male genital segment is testaceous.

Peritropis nigripennis n. sp.

Lata ovalis, nigra, capite (apice fuscescente excepto), callis pronoti medioque prosterni ferrugineis, vittis quinque basalibus pronoti fulvidis, summo apice scutelli et macula minuta angulorum basarium apicalisque cunei albidis, membrana fumosa, sparsim albido-guttulata, vagina feminae pallescente; supra squamulis seu atomis minutissimus albidis adspersa; antennae et rostrum fusco-picea; coxae nigrae, summo apice posteriorum cum trochanteribus dilute testacea (ceteræ partes pedum desunt). Caput pronoto paullo brevius, parte anteculari a supero visa oculo aequæ longa, vertice oculo duplo latiore, sulco ejus medio angusto sed distinctissimo, usque ad basin clypei extenso; articulis duobis primis antennarum glabris, primo circiter dimidio suo apicem capitis superante, latitudini verticis fere aequæ longo, rostro coxas posticas nonnihil superante. Pronotum capite duplo latius, transversim modice convexum, lateribus subrectis, angulis humeralibus rotundatis, basi parum sinuata, callis sulco tenuissimo separatis, antice collare haud tegen-