

and moderately incrassated from base to apex, second joint about twice the length of first, third a little longer than second, rostrum reaching base of ultimate ventral segment, pilose beneath, more densely and longly so on the two last joints, first and second joints of equal length, third somewhat shorter than the preceding ones together, fourth half the length of third. Pronotum rather shortly erectly palely pilose, lateral margins straight, basal margin a little rounded, the disc thickly punctate excepting the calli which are separated by a narrow linear impression. Scutellum strongly convex, sparsely punctate, pilose the same as pronotum. Corium and clavus rather thickly punctulate, suberectly pubescent, cuneus about twice longer than broad. Venter palely pilose, male genital segment at apex on each side with a rather deep and narrow sinuosity. Fore femora narrowed toward apex. First tarsal joint considerably longer than the others together. Length, ♂ 4,5 mm., incl. tegm. 6,5 mm.

Peru (Callanga). — Mus. Stockholm.

Somewhat allied to the preceding species, but with upwardly much more produced eyes, shorter antennae and legs, quite different colour-markings, etc.

Cylapus festinabundus n. sp. — A little shining, fuscous, head (except vertex), broad lateral borders of pronotum, costal margin of corium, propleurae, middle of meso- and metasternum, orificia and posterior margin of metapleura luteous, an oblong submedian spot to clavus and a large irregular area occupying the greatest part of posterior half of corium opaque, velvety fuscous-black, a spot at basal angles of scutellum and its extreme apex, veins of corium and clavus (except here and there where they run through the opaque spots), a small subcostal spot just behind middle of corium, apical angle of corium, and base of cuneus whitish, membrane with an oblong subhyaline spot at apex of cuneus, venter pale greenish, the base at the sides, the lateral margins and the male genital segment dark castaneous; antennae fuscous, first joint dingy testaceous, apex of second joint narrowly whitish; rostrum and legs brownish testaceous, hind femora with two or three rows of small fuscous spots, all tibiae fuscous-black toward apex,