in the apical half with a white annulation which is much broader in the hind tibiae. Head seen from before as high as broad, eves not much raised above vertex, their vertical diameter a little shorter than genae, vertex about two-thirds broader than an eye, first joint of antennae as long as breadth of vertex, narrowed at base, its exterior margin straight, the interior margin rounded, more so toward base. second joint about four times as long as first, third subequal in length to second, rostrum reaching base of venter, first two joints of equal length, third slightly shorter than second and distinctly longer than fourth. Pronotum with arcuately sinuate lateral margins, basal margin broadly and rather deeply bisinuate, the disc thickly punctate, excepting the calli which are separated by a posteriorly foveately deepened impression. Scutellum rather flat, broadly subimpressed in its anterior half, punctulate, rather longly erectly pilose. Corium and clavus sparsely erectly pilose, thickly punctulate except in the opaque areas, cuneus more than twice as long as broad. Venter pubescent, male genital claspers slender, sickle-shaped. Fore femora narrowed toward apex. First tarsal joint shorter than the others together. Length, 24.3 mm., incl. tegm. 5,5 mm.

Peru (Callanga) — Mus. Stockholm.

This and the following species do not seem to be nearly allied to any described form.

Cylapus labeculosus n. sp. — Above (except vertex and neck) opaque, rather sparsely pilose, beneath slightly shining, subglabrous. Head pale yellow variegated with fuscous, its blackish shining neck with two pale yellow vittae; seen from before the head is broader than high, eyes very large and prominent, but moderately raised above the vertex, their vertical diameter slightly longer than genae, vertex distinctly narrower than an eye, the clypeal prominenence distinct, the keel separating juga from genae rather strongly curved, the curve turning its convexity backwards, antennae fuscousblack, first joint dingy ochraceous with whitish base, as long as width of vertex, subparallel from apex to middle, then narrowed toward base, its exterior margin straight, interior margin slightly rounded, second joint paler toward base, considerably longer than second, rostrum fuscous-black,