

joint of hind tarsi much longer than the others, third longer than first, claws somewhat beyond middle strongly curved, pseudarolia reaching the curvature of the claws, rather broad, entirely connected with the claws, their free inferior margin incrassated, much thicker than the laminate part of the pseudarolia, which is transversely striated.

I am unable to find any described form allied to this genus.

*Pliniella sacerdotula* n. sp. — Sparingly clothed with long white semirecumbent hairs; white, underneath slightly tinted with greenish-yellow, head with the following black markings: a lateral spot to the neck extended downward in the form of a vertical fascia emitting from its middle a short vitta backward, a median vitta to vertex continued through the frons and clypeus, the narrow transverse impression between vertex and frons, a narrowly triangular spot on each side between this impression and antennal base, a vitta on the juga extended from base of antennae downward and joined to an other vitta extended from the lower angle of the eyes through the lora to apex of peristomium, these two vittae forming together a narrowly Y-shaped figure, pronotum, mesoscutum and scutellum velvety black, pronotum in its apical half with two vittae near each other and a percurrent sublateral vitta white, corium with an irregularly triangular, anteriorly sinuate dark fuscous spot at the interior apical angle, membrane dark smoky fuscous, somewhat paler in the interior half and with three rather large hyaline spots, an oblong irregular one at the basal margin, an oblong one at the interior margin, and a round one at apex, the basal hyaline spot extended from the exterior basal angle of the membrane to the inmost third of the interior cell, but interrupted by a vitta emitted from the fuscous area to the margin of the cuneus between the two outer thirds of the interior cell, and by a fuscous indentation behind the exterior cell, the incrassated basal part of the anal vein fuscous; antennae white, first joint, apical fourth of second, and apex of third black (fourth lacking); rostrum and legs white, slightly tinted with yellowish. Head a little more than half the length of pronotum, antennae linear, second joint about