

"swept from coarse grasses in a swampy opening in the pine forest." Not recorded elsewhere.

703 (—). *PHYTOCORIS UNIFORMIS* Knight, 1923, 643.

Elongate-oblong. Uniform pale brownish-yellow, rather thickly clothed with golden-yellow recumbent pubescence, intermixed with suberect pale brown hairs, eyes pale brown; femora and first antennal pale yellow, rather thickly flecked with reddish-brown spots and dots; second antennal greenish-yellow, fuscous toward apex; membrane uniformly pale dusky translucent; hind femora with a large yellow spot or ring on apical fourth. Basal joint of antennæ four-fifths as long as width of head, 2 two and two-fifths longer than 1, apical segments missing. Length, 5.2—5.5 mm.

Long Island, N. Y., Sept. 8 (*Davis*). Rocky Mountain, N. Car., Aug. 25 (*Brimley*). Recorded only from Massachusetts and New York.

704 (—). *PHYTOCORIS MEGALOPSIS* sp. nov.

Elongate, subparallel. Color above a nearly uniform fuscous-brown, thickly clothed with fine whitish appressed silky hairs and suberect dusky ones; vertex and lower half of face dull yellow; loræ tinged with reddish; eyes deep black, very large in male; collar and hind edge of pronotum whitish; costal margins of elytra and cuneus with alternate minute fuscous and white spots; inner margin of cuneus with two submedian black dots; membrane pale translucent with an iridescent sheen; femora and joint 1 of antennæ dull yellow, thickly flecked with reddish-brown spots and dots, third and fourth antennals fuscous, the former with extreme base pale; tip of beak, tarsal claws and tibial spines fuscous-brown; under surface pale reddish-brown, middle of prosternum and front coxæ ivory-white. Joint 1 of antennæ as long as width of vertex and one eye, 2 three and a half times as long as 1, 3 two-fifths the length of 2, one-third longer than 4. Sides of elytra straight and parallel to base of cuneus, thence feebly converging into the conjointly broadly rounded tips. Length to end of membrane, 5.5 mm.

Dunedin, Fla., April 20, one male taken at electric light. Due to the very large eyes the width of interocular area is but slightly more than one-half that of *uniformis*. The large coal-black eyes, uniform hue of second antennal and alternate maculation of costal margin easily separate this from the other members of Group II.

705 (—). *PHYTOCORIS DIVERSUS* Knight, 1920, 60.

Elongate, subparallel, more slender than *consersipes*. Head, pronotum and under surface greenish-yellow, the sides of tylus, margins of cheeks, oblique striæ on front, sternum and sides of ventrals usually tinged with reddish-brown; femora and tibiæ reddish-brown, mottled with paler. Joint 1 of antennæ greenish-yellow tinged with brownish, mottled with paler on inner side, one-fourth shorter than width of head;