

1 one-fourth longer than width of head across eyes, 2 two and one-fourth times longer than 1, 3 nearly three-fifths the length of 2, about one-third longer than 4. Beak reaching fifth ventral. Length, 5—5.2 mm.

Described from Connecticut. Known also from Alabama.

731 (—). *PHYTOCORIS ANGSTIFRONS* Knight Ms.

Elongate, subparallel. Pale dull yellow; elytra thickly marked with minute whitish tufts of short hairs; vertex, collar and calli whitish-yellow; basal half and sides of pronotum often with a brownish tinge; embolium, tip and inner margin of cuneus with small fuscous dots or spots; membrane pale hyaline, larger cell and two vague spots behind it fuscous, veins whitish; legs pale straw-yellow, apical half of femora pale fuscous, enclosing scattered white dots, hind femora with preapical pale ring; under surface dull yellow, a fuscous stripe along sides of ventrals. Joint 1 of antennæ pale yellow, flecked with brown, one-fifth longer than width of head across eyes; 2 dull yellow, three times as long as 1; 3 fuscous, narrowly paler at base, two-fifths the length of 2; 4 fuscous, as long as 3. Length, 4.5—5.2 mm.

Dunedin and Istokpoga, Fla., Feb. 22—April 20. Beaten in some numbers from bunches of Spanish moss where it was hibernating. Taken by Van Duzee, May 1, at Sevenoaks, Fla. Known only from that State.

732 (949). *PHYTOCORIS INFUSCATUS* Reuter, 1909, 20.

Elongate, subparallel. Head brownish-yellow flecked with red; pronotum with collar and middle of disk dull yellow tinged with reddish, sides and basal third fusco-reddish; elytra with clavus and inner half of corium dark reddish sparsely and vaguely mottled with small yellow spots; embolium and cuneus more distinctly reddish, the latter with base darker; corium with a paler spot on apex; membrane dusky, faintly marbled with pale blotches; antennæ yellowish, the outer joints with a dusky tinge; front and middle legs pale yellow, their femora tinged with reddish; hind femora except basal third, dark fusco-reddish, flecked with small yellow spots; tarsi fuscous. Joint 1 of antennæ one-fifth longer than pronotum, 2 two and three-fourth times longer than 1, 3 about one-half the length of 2, one-third longer than 4. Beak almost reaching base of genital. Length, 5.3—5.7 mm.

Marion Co., Ind., Sept. 25 (*W. S. B.*); one beaten from willow. Aqueduct, N. Y., July 27 (*Davis*). Ranges from Ontario and Massachusetts west to Michigan and southwest to North Carolina. Breeds on hickory and walnut.

733 (—). *PHYTOCORIS OLSENI* Knight, 1923, 647.

Elongate, subparallel. Head pale yellow; loræ, cheeks, middle of tylus median line and sides of base of vertex and oblique lines each side of front, orange-red; pronotum pale brownish-yellow, sides tinged with brownish, calli with a mark in front and a spot behind inner half of