duced below the eyes; vertex in male much narrower than eyes, its base not carinate; pronotum trapezoidal, its apex only onethird the width of base, disk finely, densely and evenly punctate, its hind portion strongly convex, declivent forward, collar narrow but distinct, side margins obtusely carinate, calli small, the areas in front of and between them finely punctate, hind margin broadly rounded; scutellum convex, as long as commissure, obtusely carinate at middle, distinctly punctate; elytra entire, smooth, opaque, slightly surpassing abdomen, clavus and cuneus both strongly declivent. Three neotropical species are known, one from our territory.

821 (---). CYRTOCAPSIDEA VARIEGATA sp. nov.

Short, oval. Head brown, both it and front half of pronotum thickly clothed with golden-yellow scale-like hairs; pronotum fuscous-brown, its sides and basal angles dull yellow; scutellum reddish-brown, its carina and apex dull yellow; clavus fuscous, paler at base and apex; corium fuscous along claval suture, outer half dull yellow; embolium fuscous, vaguely flecked with small yellowish spots; cuneus reddish-fuscous, its tip and inner margin pale yellow; membrane dusky hyaline, basal half of cells fuscous, veins and outer edges pale; legs pale yellow, femora tinged with fuscous; under surface fuscous-brown, middle of ventrals and genital paler. Joints 1 and 2 of antennæ straw-yellow, 1 slightly longer than width of vertex, 2 distinctly thicker toward apex, three and a half times longer than 1; 3 and 4 fuscous, united subequal in length to 2. Elytra and base of scutellum thinly clothed with yellow, scale-like hairs. Length, 2.4 mm.

R. P. Park, Fla., March 17; one male swept from roadside herbage. Easily known from our other species of Capsaria by its short oval form, very large eyes, opaque impunctate elytra and floccose yellow scales on head and front half of pronotum.

XXII. NEOBORUS Distant, 1884, 276.

Oblong or oblong-oval shining species having the head immersed in thorax to eyes, its front nearly vertical; eyes very prominent, compressed, overlapping front angles of pronotum; vertex subdepressed, its base margined; antennæ shorter than body, joint 1 glabrous, but little thicker than 2; pronotum trapezoidal, its sides straight, their margins carinate, calli smooth, prominent, the area between and in front of them punctate, hind portion convex, declivent forward, coarsely punctate, hind angles obtusely rounded, basal margin subtruncate; scutellum feebly convex, sparsely irregularly punctate;