

*Fulgentius mandarinus*, sp. n.

Black; antennæ, eyes, legs, and membrane piceous; anterior margin of pronotum, first joint of antennæ (excluding apex and base of third joint) and apices of femora ochraceous; corium with a transverse fascia before middle and between clavus and lateral margin, and about basal half of cuneus, greyish white; body beneath black, imperfectly seen in carded specimen, but apparently with a greyish spot near posterior coxæ; body above very finely and obscurely pilose.

Long. 8 mm.

*Hab.* China; Namoa Islands (*J. J. Walker*, Brit. Mus.).

## NICHOMACHUS, gen. nov.

Moderately elongate. Head broad, including eyes much wider than anterior margin of pronotum, narrowed and moderately deflexed in front of the prominent and exerted eyes, lateral margin sinuate, disk strongly longitudinally sulcate; antennæ with the first joint short, shorter than antero-ocular portion of head, second and third joints longest and subequal in length, fourth shorter but longer than first; rostrum reaching the posterior coxæ; pronotum strongly constricted at about one third from anterior margin, forming a distinct narrow anterior lobe, posterior lobe tumid, about twice as long and much broader than the anterior lobe; scutellum very strongly conically gibbous and longly though sparingly pilose; corium (excluding cuneus) a little shorter than the abdomen, its lateral margins sinuate, broadest at the area of the interior angle, cuneus longer than broad; membrane thickly and finely reticulate, with a single, narrow, short, lateral cell.

Allied to *Systemonotus*, from which it differs by the broader head, larger and exerted eyes, conically raised scutellum, &c. But for the longitudinally impressed head might be located in the *Pilophoraria*.

*Nichomachus Sloggetti*, sp. n.

Cinnamon-brown; eyes, scutellum, base and apical margin of corium, cuneus, disks of meso- and metasterna, and abdomen beneath black; an oblique transverse fascia in basal black area of corium, a transverse fascia to clavus beyond middle, and a basal fascia to cuneus white; antennæ (excluding basal joint), posterior lobe of pronotum, and apices of femora infuscated; membrane shining brownish ochraceous; two transverse subbasal fasciæ to abdomen beneath pale