

*Note*.—In this division *Miraria* and near the genus *Pantilius* I place *Pæas Reuteri*, Dist. (Biol. Centr.-Am., Rhyn. i. p. 428, tab. xxxvii. fig. 5),—head distinctly sulcated; and for the same reason *Jacchinus tabascoensis*, Dist. (*loc. cit.* p. 430, tab. xxxvii. fig. 10).

#### Division CYLAPARIA.

*Valdasaria*, Dist. Biol. Centr.-Amer., Rhynch. i. p. 242 (1883).

*Monaltonionaria*, Reut. Ann. Soc. Ent. Fr. lxi. p. 398 (1892).

*Eucerocharia*, Kirk. J. Bomb. Nat. Hist. Soc. 1902, p. 294.

*Cylaparia*, Kirk. Wien. ent. Zeit. xxii. p. 13 (1903).

The name of this division was originally founded on that of the neotropical genus *Valdasus*, Stål; this having since been proved to be but a synonym of *Cylapus*, Say, it is necessary to alter the divisional name as above.

#### ARGENIS, gen. nov.

Head broad, not horizontally produced in front of eyes, which touch but exceed the width of the anterior margin of the pronotum, distinctly longitudinally centrally impressed, or very finely sulcate; antennæ with the first joint longer than head, but shorter than pronotum, second joint almost twice as long as first, third joint about one third shorter than second; eyes large, globose; pronotum coarsely punctate, transversely constricted before middle, posterior lobe tumid, centrally very obscurely carinate, posterior angles subprominent and a little tuberculous; scutellum triangular, the apex somewhat acute; corium long, cuneus passing apex of abdomen; membrane somewhat small; body beneath obscurely seen, owing to typical specimen being carded.

#### *Argenis incisuratus*.

*Capsus incisuratus*, Walk. Cat. Het. vi. p. 121. n. 282 (1873).

*Hab.* Ceylon.

Walker's very inadequate description of this species contains the erroneous statement:—"Prothorax with no transverse furrow."

#### Genus SYSINAS.

#### *Sysinas tibialis*.

*Capsus tibialis*, Walk. Cat. Het. vi. p. 109. n. 245 (1873).

*Resthenia tibialis*, Atkins. Cat. Capsidæ, p. 61 (1890).