

Division PHYTOCORARIA.

Genus PARACALOCORIS.

Paracalocoris Barretti, sp. n.

Purplish brown; head, antennæ (excluding basal joint), lateral margins and a broad central fascia (attenuated posteriorly) to pronotum, basal angles of scutellum, a very small marginal spot near apex of corium, a marginal spot to cuneus, body beneath, rostrum, and legs pale ochraceous; apices of second and third joints of antennæ and apices of tibiæ purplish red; pronotum with two discal black spots; first joint of antennæ incrassate and pilose, second joint distinctly incrassate towards apex, about half as long again as first; pronotum transversely rugulose; membrane very pale fuscous with the veins darker.

Long. $5\frac{1}{2}$ mm.

Hab. Cape Colony: King William's Town (*Miss Barrett*, Brit. Mus.).

Division CAPSARIA.

Genus LYGUS.

Lygus Schonlandi, sp. n.

Ochraceous; hemelytra somewhat longly pilose; apex of second joint of antennæ black (remaining joints mutilated); basal area of pronotum, two central longitudinal fasciæ to scutellum, inner area and two lateral spots (one before middle, the other at apex) to corium, and a spot at apex of cuneus piceous; basal and inner margins of cuneus generally distinctly narrowly sanguineous; membrane fuscous with paler mottlings; body beneath and legs pale ochraceous; mesosternum, a lateral spot to metasternum, base of posterior tibiæ, and apices of tarsi black; apical halves of posterior femora castaneous with broad fuscous annulations; rostrum reaching the intermediate coxæ; pronotum finely and obscurely punctate; first joint of antennæ slightly thickened, second joint a little more than twice the length of first.

Long. 4 to $4\frac{1}{3}$ mm.

Hab. Cape Colony: Grahamstown (Albany and Brit. Mus.). Natal: Durban (*Marshall*).