

and corium a little more finely and obscurely punctate; cuneus sanguineous, its outer area and apex black.

Long. 5 mm.

*Hab.* Transvaal: Zoutpansberg (*Junod*, Brit. Mus.).

*Camptobrochis capensis*, sp. n.

Reddish ochraceous; head and scutellum black, the last with a central reddish-ochraceous fascia, which does not reach the base; antennæ, lateral margins of corium, tibiæ, and tarsi pale ochraceous; apices of tarsi black; antennæ somewhat slender, second joint more than twice the length of first; pronotum distinctly punctate, anterior and posterior margins narrowly ochraceous, the last linearly transversely black near lateral angles; scutellum and corium more finely and obscurely punctate than pronotum; corium and clavus somewhat longly pilose; membrane fuscous, with paler mottlings.

Long. 4 mm.

*Hab.* Cape Colony: Grahamstown (Albany and Brit. Mus.).

Division BRYOCORARIA.

Genus TENTHECORIS.

*Tenthecoris*, Scott, Ent. Month. Mag. xxxiii. p. 65 (1886).

Type, *T. bicolor*, Scott (Brit. Mus.).

This genus is very closely allied to *Ecritotarsus*, Stål. It is described as an orchid pest, as is also *Ecritotarsus exitiosus*, Dist., and *E. orchidearum*, Reut. *T. bicolor* is very closely allied by description to Reuter's species; Scott describes the first and second joints of the antennæ as red, but in one of his typical specimens the apex of the first joint and the whole of the second joint are distinctly black.

Division — ?

Genus FUNDANIUS.

*Fundanius alternus*.

*Capsus alternus*, Walk. Cat. Het. vi. p. 111. n. 252 (1873).

*Resthenia alternus*, Atkins. Cat. Capsidæ, p. 57 (1890).

Division — ?

Genus ARMACHANUS.

*Armachanus spicatus*, sp. n.

Uniform pale cinnamon-brown; a discal, transverse, pale greyish line across apex of clavus, and an oblique line of the same colour crossing corium near middle; a prominent black