

C. *Pronotum with two prominent black spots.*1422. *Lygus catullus*, sp. n.

Head and pronotum pale ochraceous; eyes, extreme apex of head, and two spots on anterior area of pronotum black; scutellum and corium dark shining fuscous; two small basal spots and apex to scutellum, and basal and apical marginal spots to corium pale ochraceous; cuneus pale ochraceous with an inner fuscous spot; membrane very pale fuliginous, base irrorated with pale flavescent and fuscous; body beneath and legs pale ochraceous; a lateral spot to meso- and metasterna, and some transverse fasciæ and the apex to abdomen dark fuscous; pronotum very obsoletely finely punctate; scutellum (excluding apex) finely transversely striate; corium with some scattered pale flavescent hairs; antennæ pale ochraceous, extreme apex of first, apical half of second, and the third joint (excluding base) fuscous.

Length 4 millim.

*Hab.* Tenasserim; Myitta (*Doherty*).

D. *Pronotum with a central dark longitudinal fascia.*1423. *Lygus biseratensis*, *Dist. Fasciculi Malayenses*, *Zool.* vol. i, p. 269, pl. xvi, ff. 11, 11 a (1903).

Pale obscure ochraceous, finely sericeous; apices of first, second, and fourth joints of antennæ, eyes, a submarginal anterior line and a central longitudinal fascia to pronotum black; cuneus fuscous-brown; first and fourth, and second and third joints of antennæ subequal in length.

*Var. a.* Scutellum and corium pale purplish-brown, the lateral margins of the last pale ochraceous.

*Var. b.* Scutellum and corium dark castaneous, the lateral margins of the last narrowly pale ochraceous; femora moderately infuscated.

Length 7 millim.

*Hab.* Ceylon; Kandy (*Green*). Burma; Mandalay (*Fea*).—Siamese Malay States.

E. *Species unknown to the writer.*1424. *Lygus bengalicus*, *Reut. Ent. Tidskr.* v, p. 195 (1884).

Green, somewhat palely pubescent; head, antennæ, apices of femora, tibiæ, and tarsi more or less reddish-brown; apex of clypeus piceous; antennæ with about a third of basal joint and apex of second somewhat fuscous, frequently in the ♂ with the second joint totally fuscous; pronotum piceous, with a large anterior rounded greenish-livid spot in ♂, or sordidly greenish, with two submarginal broad stripes, sometimes broadly dilated posteriorly and before the basal margin confluent in ♀; hemelytra