

with a fuscous spot at the interior apical angle of corium; membrane fuliginous, the cells immaculate or with their apices palely infuscated; a hyaline groove at apex of cuneus in ♂, or two semihyaline spots before the middle of the external margin in the ♀; apices of tarsi darkly fuscous; rostrum reaching posterior coxæ; hemelytra densely punctate.

Length  $3\frac{1}{2}$  to 4 millim.

*Hab.* Bengal (*vide Reuter*).

I have not seen this species.

1425. *Lygus obtusus*, *Reut. Ent. Tidskr.* v, p. 196 (1884).

Dark green, somewhat glabrous, slightly punctate; eyes fuscous; clypeus, a band on the pleura of the prothorax, an apical stricture and medial basal margin of pronotum, a medial band to scutellum, all the claval sutures (interiorly broad), an apical spot to corium, apical margin to cuneus (interrupted medially), a common arcuated fascia to the apices of the cells and two spots on the external edge of membrane, another a little behind apex of cuneus and another situate behind middle fuscous; legs dark green, unspotted, tibiæ darkly spinulose, last joint of tarsus apically dark fuscous; rostrum reaching the intermediate coxæ; hemelytra slightly punctate.

Length, ♀,  $3\frac{3}{8}$  millim.

*Hab.* Bengal (*vide Reuter*).

I have not seen this species.

Genus **PÆCILOSCYTUS.**

*Pæciloscytus*, *Fieb. Wien. ent. Monatsb.* ii, p. 311 (1859); *id. Eur. Hem.* pp. 68, 276 (1861); *Reut. Bih. Vet.-Ak. Handl.* iii, 1, p. 19 (1885); *id. Acta Soc. Faun. & Flora Fenn.* i, p. 80 (1875).

Subg. *Charagochilus*, *Fieb. Wien. ent. Monatsb.* iii, p. 309 (1859).

„ *Systratiotus*, *Dougl. & Scott, Brit. Hem.* p. 443 (1865).

Type, *P. unifasciatus*, Fabr., a Palæarctic and Nearctic species.

*Distribution.* Probably universally distributed.

Body oval, opaque or subopaque, above flavescently pilose; head nearly vertical, inner margins of eyes marked with testaceous pubescence; eyes moderately exserted; antennæ moderately slender, first joint not longer than head, second twice as long as first; rostrum reaching or only slightly passing intermediate coxæ; pronotum trapeziform, anteriorly convexly deflected, posterior margin truncate, not covering basal margin of scutellum, the disk densely punctate; cuneus very strongly fractured, its apex ferruginous or flavous.

1426. *Pæciloscytus longicornis*, *Reut. (Charagochilus) Ent. Tidskr.* v, p. 196 (1884).

Black, hemelytra and anterior collar to pronotum greyishly