

piceous; membrane fuliginous, with two paler spots on the lateral margins; head and anterior margin of pronotum darker luteous; rostrum reaching the posterior coxæ, its apex black.

Length 6 millim.

Hab. "North Hindostan" (*Boyes, Brit. Mus.*).

Genus LUCITANUS, nov.

Type, *L. punctatus*, Kirby.

Distribution. Ceylon.

Subovate; head somewhat long, moderately deflected before eyes, first joint of antennæ considerably longer than head and a little thickened, second joint much longer than first and thickened on apical area, third and fourth joints short, slender, third a little longer than fourth; rostrum about reaching the posterior coxæ; pronotum tumid, deflected and narrowed anteriorly, lateral margins a little rounded, basal margin slightly concave in front of scutellum and thence oblique to lateral angles; scutellum large, subtriangular, about as broad as long; clavus very long, extending almost to base of membrane; lateral margins of corium a little convex; cuneus much broader than long; legs of moderate length, posterior femora thickened; tibiæ setose.

1439. *Lucitanus punctatus*, Kirby (*Leptomerocoris*), *J. Linn. Soc., Zool.* xxiv, p. 108 (1891).




Fig. 290.—*Lucitanus punctatus*.

Pale ochraceous; head and pronotum shining; eyes, lateral margins (broken anteriorly), two anterior contiguous transverse discal spots and two rounded widely-separated central spots to pronotum, and longitudinal fasciæ to scutellum and corium, black: membrane very pale fuliginous, subhyaline, the venation and two posterior discal angulate lines black; antennæ with the apex of the second joint fuscous, third and fourth joints black, base of third stramineous.

Length 5 to 6½ millim.

Hab. Ceylon; Tangalla (*Green*).

Genus DERÆOCORIS.

Deræocoris, *Kirschb. Jahrb. V. Nat. Herz. Nassau*, Heft x, p. 208 (1855); *Stål, Hem. Afr.* iii, p. 20 (1865); *Reut. Act. Soc. Sc. Fenn.* xxiii, no. 2, p. 18 (1898), et syn.
Macrocapus, *Reut. Öfv. Finsk. Soc. Förh.* xxi, p. 55 (1879).