

fascia and faintly transversely wrinkled; clavus and pronotum pale brownish ochraceous, thickly more darkly punctate, the costal area paler and impunctate, cuneus ochraceous, the apical area piceous, somewhat longly pilose; membrane pale fuliginous; body beneath and legs ochraceous, lateral margins of sternum and abdomen and apical area of abdomen broadly piceous; other characters as in generic diagnosis.

Length 5 mm.

*Hab.* Tenasserim; Myitta (*Doherty*).

#### SOHENUS, gen. nov.

Head considerably longer than broad, tumid, eyes somewhat small and adpressed and placed about midway between base and apex, the apical area strongly deflected in front of eyes, the basal area moderately attenuated from eyes to posterior margin; antennæ rather long, first joint shorter than head, second three times as long as first, third almost as long as second and half as long again as fourth; rostrum reaching the intermediate coxæ; pronotum much broader at base than long and more than twice broader than anterior margin, three transverse impressions before anterior margin, the basal area gibbous and deflected to the first transverse impression, and the lateral margins to that impression a little convexly oblique, remaining lateral margins to apex distinctly dislocated at the remaining impressions, posterior margin angularly concave; scutellum about as long as broad at base, strongly deflected posteriorly and transversely impressed near middle, the lateral margins moderately concavely sinuate; corium (including cuneus) as long as head, pronotum, and scutellum together, the costal margin strongly concavely sinuate, apex of clavus almost reaching base of cuneus, which is a little broader than long; membrane about half the length of corium (including cuneus); legs long, posterior tibiæ about as long as scutellum and hemelytra together, posterior tarsi with the first joint a little longest.

Allied to *Armachanus*, Dist., but with the head longer, and with no apical spinous protuberance, &c.

#### *Sohenus proditus*, sp. n.

Pale cinnamomeous; eyes black; corium with a somewhat oblique white fascia crossing it and greater part of clavus a little beyond middle, apical inner area of corium somewhat infusate, membrane fuliginous; antennæ with the first, second, and fourth joints pale fuscous brown, third joint pale ochraceous; scutellum darker and a little more castaneous in