

IDATIUS, gen. nov.

Head about as long as broad, not foveately sulcate between the eyes, which project considerably beyond the anterior margins of the pronotum; antennæ with the first and second joints moderately thickened, the first distinctly shorter than head, second four times the length of first, third and fourth slender, but together not so long as second; rostrum robust, reaching the intermediate coxæ; pronotum at base twice as broad as long and about twice as broad as at anterior margin, deflected anteriorly, the lateral margins a little roundly oblique, the posterior margin subtruncate; scutellum somewhat large and triangular, moderately transversely impressed before middle; corium (excluding cuneus) considerably longer than second joint of antennæ; cuneus with the basal and outer margins subequal in length; membrane considerably passing the abdominal apex, with a distinct basal cell; posterior femora moderately thickened, posterior tibiæ spinulose.

Idatus priscillianus, sp. n.

Black; eyes black; antennæ with the basal joint brownish ochraceous, second joint black, with its basal area brownish ochraceous, third and fourth joints palely fuscous, basal area of third ochraceous; cuneus with two greyish spots near base; membrane pale fuliginous; body beneath black; legs ochraceous, posterior femora castaneous, the spinules to the posterior tibiæ and the apices of the tarsi black; pronotum and scutellum shining black, faintly transversely striate; corium obscurely pilose and finely punctate; other structural characters as in generic diagnosis.

Length 3 mm.

Hab. Bengal; Paresnath, 4000 to 4500 ft. (*Annandale*).
Simla Hills; Dharampur, 5000 ft. (*Annandale*).

SEJANUS, gen. nov.

Head about as long as broad, deeply, broadly, roundly, foveately sulcate between the eyes, which are large and project beyond the anterior margins of the pronotum; antennæ with the first and second joints moderately thickened, the second more robust than the first, which is short and about the length of the eyes, second joint about five times the length of first, third and fourth slender, third about twice as long as first, fourth as long as first; rostrum robust and reaching the posterior coxæ; pronotum twice as broad as