

Megacælum nigroscutellatum, sp. n.

Ochraceous, moderately and testaceously punctate; antennæ ochraceous, apices of the first and second joints fuscous; eyes black; pronotum with the lateral areas generally more or less testaceous, two dark spots on anterior lobe and a series of dark spots on basal margin, the whole surface distinctly punctate; scutellum with the apex broadly black; corium finely, more darkly, and testaceously punctate, the lateral marginal areas paler and sparsely maculate; cuneus finely testaceously maculate; membrane dull greyish with darker suffusions; legs pale ochraceous, the apical areas of the femora and the tibiæ with darker annulations; first joint of antennæ slightly incrassated, a little longer than head, second joint about twice as long as first; membrane passing abdominal apex.

Var.—Scutellum with two small black spots on apical area, not apically broadly black.

Long. 5 mm.

Hab. Central Districts, Upper Houadou, Gonde.

NEMESIANUS, gen. nov.

Subelongate; head with a distinct central longitudinal carination; eyes somewhat prominent, contiguous to the anterior margin of the pronotum, but moderately projecting beyond it; antennæ with the first joint moderately incrassated, about as long as head, second joint about three times as long as first, third and fourth joints short, slender, third a little longer and stouter than fourth; rostrum indistinctly seen in carded specimens; pronotum about as long as half the width at base, the basal lateral angles subprominent, the lateral margins a little concavely sinuate; scutellum triangular, about as long as broad at base; femora somewhat strongly incrassate.

Allied to *Tancredus*, Dist.

Nemesianus nigroruber, sp. n.

Head, pronotum, and scutellum dull dark sanguineous, eyes and basal area of pronotum piceous or black; antennæ with the first joint testaceous, its apex narrowly black, second joint pale testaceous, with nearly its apical half black, third and fourth joints black, their extreme bases paler; corium piceous,