FAMILY 6.—LITOSOMIDÆ.

Genus 1.—LITOSOMA, Doug. & Scott.

Species 13.—LITOSOMA BICOLOR, Doug. & Scott, pl. 2, fig. 3.

Elongate, narrow. Green, somewhat thickly clothed with fine, short, depressed, whitish hairs, intermixed with longish erect black ones.

Head—Crown generally brownish or fuscous. Antennæ, 1st and 2nd joints greenish, apex of the latter brown; 3rd and 4th fuscous. Rostrum greenish, apex piceous.

Thorax—Pronotum brownish or fuscous, lateral margins green. Scutellum convex, raised above the clavus, brownish or fuscous. Elytra—Clavus brownish or fuscous. Corium green; next the claval suture sometimes very narrowly, and between the apex of the clavus and the inner posterior angle, brownish or fuscous. Cuneus green, of the same colour as the corium. Membrane blackish, iridescent; large cell nerve blackish, lesser cell nerve yellowish. Sternum—Mesosternum blackish, anterior and posterior margins and a central line yellow. Legs greenish-yellow, clothed with fine whitish hairs. Thighs, 1st pair with a row of fine, erect, longish brown hairs. Tibiæ with long, erect, somewhat spinose brown hairs; apex brown. Tarsi, 1st and 2nd joints brownish-yellow, 3rd and claws black.

Abdomen underneath green, clothed with fine depressed whitish hairs.

Length—♂ 2, ♀ 1¾ lines.

In the Q the head and pronotum are green, and the brownish or fuscous markings on the scutellum and elytra never so decided as in the \mathcal{S} .

Not uncommon by beating furze bushes at Esher in August. The males are exceedingly active.

FAMILY 11.—ONCOTYLIDÆ.

Genus 2.—Macrocoleus, Fieb.

Species 1*.—Macrocoleus solitarius, Meyer, pl. 2, fig. 4.

Capsus solitarius, *Meyer*, Caps., 83, 62, t. 5, fig. 4 (1843).

Capsus seladonicus, Kirschb., Caps., 80, 103 (1855).

MACROCOLEUS SOLITARIUS, Fieb., Europ. Hem., 321, 8.

Elongate, grey-greenish or grey-yellowish, clothed with stout erect black hairs; the nerves of the elytra naked, pale.

- Head elongate, pointed, greenish-yellow. Crown convex. Antennæ dingy green, clothed with short black hairs; 2nd joint at the apex narrowly, and 3rd and 4th, brownish. Rostrum brown, apex piceous.
- Thorax—Pronotum greenish-yellow; callosities not divided, their hinder margin emarginate; posterior margin at least three times as wide as the anterior; disc convex. Scutellum raised above the clavus, with a transverse channel