

keels; side margins of the *pronotum* and anterior margin of the *elytra* wide, with four rows of meshes, the circumference of the former broadly rounded anteriorly.

In structure, marking, colour, and size, this species is very like *M. ampliata*, from which it differs in the following respects. The antennæ are perceptibly thicker, and much shorter, the spines on the head longer and sharper, the side margins of the *pronotum* more rounded at the front, the curve being continued regularly to the hood (in *ampliata* the margin *projects* anteriorly in an obtuse angle, and then goes in an oblique straight line to the hood).

Two specimens taken by Mr. Wollaston, but the place and date of capture are not recorded.

Section 9.—CAPSINA.

FAMILY 3.—MIRIDÆ.

Genus 2A.—TERATOCORIS, Fieb.

Species 2.—TERATOCORIS SAUNDERSI, n. sp.

Bright green, shining, sparingly clothed with short, sub-erect, yellow hairs.

♂. *Head*—With a black central line extending throughout its entire length, widest at the posterior margin; round the insertion of the antennæ narrowly black. *Antennæ*, 1st joint green, clothed with short dark hairs; base with a narrow black ring, apex more or less pitchy-red; 2nd brownish-pink, apex brown; 3rd and 4th pitchy-black. *Eyes* reddish-black. *Rostrum* greenish-yellow, apex black.

Thorax—*Pronotum* green, with a black central line extending throughout its entire length; at the anterior angles a short, somewhat oblique black streak reaching to the callosities, the latter with a slight fovea near the centre; collar and posterior portion of the disc finely shagreened. *Scutellum* green, with a more or less distinct short black streak between the base and the transverse channel; at the basal angles a deep fovea. *Elytra* green, as long as the abdomen, finely shagreened. *Clavus*, inner margin very narrowly black. *Membrane* with only one cell, pale fuscous with a purple iridescence. *Cell* green, almost entirely subcoriaceous and finely shagreened; cell nerve green. *Sternum* green. *Prosternum*, on the sides towards the front, with a short, cuneate, black patch. *Legs* green, clothed with short, sub-depressed, dark hairs, *Thighs*, at the apex, more or less inclined to reddish; 1st and 2nd pairs, on the underside, with a row of erect hairs. *Tibiae* greenish or greenish-yellow, thickly clothed with short, sub-depressed, brownish-yellow hairs; 2nd and 3rd pairs frequently inclined to brownish-pink or yellow at the apex. *Tarsi* brownish-yellow, apex of 3rd joint black. *Claws* brown.

Abdomen green underneath.