

- ♀. Developed form. *Head* and *pronotum* as in the ♂, the central line of the head not so distinct. *Scutellum* without a central line; apical portion transversely wrinkled. *Elytra* longer than the abdomen. *Membrane* with two cells, the lesser one very narrow, and almost forming an isosceles triangle. All the other characters as in the ♂.
- ♀. Undeveloped form. *Head*, *pronotum*, and *scutellum* without the black central line; posterior margin of the callosities blackish. *Elytra* shorter than the abdomen. *Cuneus* not distinct from the corium. All the other characters as in the developed form. Length ♂ 2—2½; ♀ 2½—3 lines.

Most nearly allied to *T. antennatus*, Boh. (Fieb. Europ. Hem. 246, 1), but the absence of the streak along the margin of the abdomen, and the blood-red hinder tibiæ, will at once enable any one to separate them.

We have much pleasure in naming the species after Mr. Edward Saunders, its captor, who took a few examples at Deal, by sweeping among rushes, &c., at the end of June and beginning of July. He has also an undeveloped ♀, taken near Aberdeen.

Genus 3.—LOPOMORPHUS.

We now believe that the insect described in the "British Hemiptera," page 224, 1, as *Lopomorphus carinatus*, is only a small and curious variety of *L. ferrugatus*, to which the description must also apply.

FAMILY 4.—PHYTOCORIDÆ.

Genus 2.—PHYTOCORIS, Fall.

Species 2A.—PHYTOCORIS MARMORATUS, n. sp.

Pale green, with large irregular black patches, sometimes almost covering the entire elytra; at others, having somewhat of a banded appearance; clothed with depressed white hairs, slightly curled, and disposed in a confused manner, and interspersed with sub-erect black ones.

Head—pale yellowish or greenish-white at the posterior margin, and adjoining ear eye a small piceous or blackish spot. *Antennæ* black, as long as the body; 1st joint a little more than half the length of the second, with one or two small, somewhat round white spots towards the base, and two or three oblong ones towards the apex on the upper side, and a few long, erect, black hairs; apex slightly piceous, 2nd with a narrow white ring at the base, and another, whitish or brownish-white, of about the same size beyond the middle; 3rd about two-thirds the length of the second, base narrowly white; 4th shorter than the 1st; below the eyes and beyond the side-lobes of the face a black streak. *Rostrum* pale-yellowish or greenish-white, apex piceous.