

**Hadronema princeps** Uhl.

*Hadronema princeps* Uhler, Proc. Calif. Acad. Sci., ser. 2, vol. IV, 1894.

A much narrower form than the other species with usually a distinct light mark on the cuneus. Occurs throughout the West.

**Hadronema picta** Uhl.

*Hadronema picta* Uhler, Colo. Exp. Sta. Bull. No. 31, p. 31, 1895.

Separable from the other species in having the second and third joints of the antennæ equal in length. Recorded as occurring in Colorado and Dakota.

**Hadronema festiva** Van D.

*Hadronema festiva* Van Duzee, Trans. Am. Ent. Soc., vol. XXXVI, No. 2, p. 80, 1910.

The only species of the genus having the third joint of the antennæ longer than the second. Also readily distinguishable by the radically different colour markings.

Mr. Van Duzee records its occurrence in New Mexico.

**Hadronema splendida**, n. sp.

Resembles *festiva* Van D. Head small, vertical, black with light markings next to eyes, near base of head, on the centre of the face, and below the antennæ. Bucculæ bordered with white. Antennæ black, third joint noticeably shorter than second. Pronotum dark gray or black, callosities black, anterior border of pronotum light. Scutellum red. Elytra greenish white, smoky at middle of corium, clavus smoky along inner margin, cuneus white except smoky at apex. Costal border of elytra whitish. Membrane smoky. Abdomen dark beneath but light at the middle. Femora bright orange red, tibia and tarsi black. Size 3.8 mm. long, 1.2 mm. wide.

Described from a female collected at Albuquerque, N.M., Aug., 1909, now in the U. S. National Museum.

The peculiar colour markings readily separate this species from all others.