

more slender near base; III, .33 mm.; IV, .25 mm.; uniformly pale yellowish brown; clothed with fine yellowish pubescence. Rostrum: length .71 mm., reaching posterior margin of intermediate coxae, yellow, apex brown. Pronotum: length .42 mm., width at base .96 mm.

Clothed with simple, coarse, yellowish to golden pubescence; uniformly pale yellowish to brownish above and below; membrane dusky, veins sometimes paler.

Female. Length 2.6 mm., width 1.1 mm. Head: width .56 mm., vertex .32 mm. Antennae: segment I, length .13 mm.; II, .58 mm., scarcely equal in thickness to segment I, much thinner on basal half; III, .31 mm.; IV, .25 mm. Pronotum: length .42 mm., width at base .92 mm. Coloration usually a paler yellowish brown than the male, pubescence very similar.

Holotype: male, May 25, 1931, Corinth, Miss. (H. G. Johnston); author's collection.

Allotype: taken with the type.

Paratypes: 10 males and females taken with the types on wild grape (*Vitis* sp.). 6 males and females May 15, 1931, Natchez, Miss. (H. G. Johnston), also on wild grape.

***Neoborus osmanthicola* n. sp.**

The coloration suggestive of *canadensis* V. D., but differs in the broadly pale margins of scutellum, nearly glabrous surface of the dorsum, the less prominent eyes and shorter rostrum.

Male. Length 4.6 mm., width 1.9 mm. Head: width 1.01 mm., vertex .42 mm.; vertex and frons sharply declivent; yellowish, tylus brownish black with few scattered, yellowish hairs. Rostrum, length 1.09 mm., scarcely attaining the posterior margin of mesosternum, yellowish, apex black. Antennae: segment I, length .54 mm., brownish black, nearly glabrous; II, 1.36 mm., yellowish brown becoming brownish black basally; III, .52 mm., fuscous; IV, .33 mm., fuscous. Pronotum: length .88 mm., width at base 1.56 mm., lateral margins straight, but scarcely carinate.

Dorsum nearly glabrous, shining, a few rather conspicuous hairs on embolium, outer margin of corium and cuneus; short, inconspicuous hairs in punctures on clavus, inner apical margin of corium, and hind margin of pronotum. Coloration yellowish brown and marked with reddish brown; clavus reddish brown, paler along apex of claval vein; corium on apical one-third reddish brown, anal ridge and extending along inner apical angle fuscous to black; apex of embolium pale yellowish; cuneus yellowish, apical one-half dark red. Scutellum yellowish brown, side margins yellow, impunctate. Membrane